

EFFECTS OF LAND USE CHANGE ON FIRE, VEGETATION AND  
WILDLIFE DYNAMICS IN ARID ECOSYSTEMS OF SOUTHERN  
RUSSIA

by

Maxim Dubinin

A dissertation submitted in partial fulfillment  
of the requirements for the degree of

Doctor of Philosophy  
(Forestry)

at the

UNIVERSITY OF WISCONSIN-MADISON

2010

EFFECTS OF LAND USE CHANGE ON FIRE, VEGETATION AND  
WILDLIFE DYNAMICS IN ARID ECOSYSTEMS OF SOUTHERN  
RUSSIA

submitted to the Graduate School of the  
University of Wisconsin-Madison  
in partial fulfillment of the requirements for the  
degree of Doctor of Philosophy

by  
Maxim Dubinin

Date of final oral examination: November 23, 2010

Month and year degree to be awarded: December 2010

The dissertation is approved by the following members of the Final Oral Committee:

Radeloff, Volker C., Associate Professor, Forest and Wildlife Ecology

Turner, Monica G., Professor, Zoology

Jackson, Randall D., Associate Professor, Agronomy

Townsend, Phil, Professor, Forest and Wildlife Ecology

Van Deelen, Tim R., Assistant Professor, Forest and Wildlife Ecology

# EFFECTS OF LAND USE CHANGE ON FIRE, VEGETATION AND WILDLIFE DYNAMICS IN ARID ECOSYSTEMS OF SOUTHERN RUSSIA

**Maxim Dubinin**

**Under supervision of Associate Professor Volker C. Radeloff**

**At the University of Wisconsin-Madison**

Human land use is widely acknowledged to be the most influential disturbance agent in ecosystems around the globe. Changes in the disturbance regime profoundly affect components of natural and coupled socio-ecological systems such as land cover, ecosystem processes, and biodiversity. Excessive increases in disturbance may lead to catastrophic shifts in the composition and function of communities, as critical thresholds of disturbance are reached.

Decreases in land use intensity are less common, and less studied, but no less important and provide opportunity to understand ecosystem recovery.

The overarching goal of my dissertation was to study the effects of decreasing land use intensity on fire, vegetation, and wildlife dynamics in arid ecosystems. I chose the south of European Russia as my study area because the collapse of the Soviet Union here led to rapid changes in the main land use – sheep grazing – which caused substantial environmental changes. I used a variety of remote sensing data and image processing methods to quantify scale and pattern of these changes from 1985 until 2007 and to link these changes with habitat selection of Saiga antelope – a species of major conservation concern.

Results revealed significant changes in ecosystem patterns and processes following the collapse of the USSR. Decrease in land use intensity led to dramatic increase in fires from almost none in the 1980s to more than 20% of the total study area burned annually in the 2000s. It took vegetation 5 to 6 years to recover the point where large scale fires could be sustained. The total area burned was driven by a number of factors, including sheep numbers, precipitation during growing season, temperatures during burning season and vegetation condition. Together with fires, vegetation composition changed substantially, switching from communities dominated by shrubs to perennial grasses. Though current habitat selection by Saiga antelope was not affected by vegetation composition at the scale of my analysis, fires played prominent role as a landscaping agent and areas that burned more often and were close to water sources were preferentially selected. In general, my dissertation shows that institutional changes may lead to both ecosystem recovery and transitions to unexpected states. Natural experiments set up by large scale socio-economic changes can provide a framework to study the effects of such changes over large areas and elucidate the direction of ecosystem development under decreasing land use conditions in other parts of the world.



## Acknowledgements

Only in full version ☺



## Table of contents

Introduction.....	1
Chapter 1: Reconstructing long time series of burned areas in arid ecosystems of Southern Russia by satellite remote sensing .....	18
Abstract .....	18
Introduction.....	20
Methods.....	24
Study area.....	24
Study period .....	25
Data .....	26
Data processing.....	28
Validation.....	30
Results.....	32
Annual burned area .....	32
Validation.....	33
Discussion .....	35
Burned area trends, vegetation, grazing, and socioeconomic changes .....	36
Satellite data limitations.....	38
Validation of the burned area estimates and the burned area mapping approach.....	39
Concluding remarks .....	40
Acknowledgements.....	41
Bibliography .....	42

## Chapter 2: Relationships between climate, livestock, and fire in arid ecosystems of Southern

Russia.....	63
Abstract .....	63
Introduction.....	64
Methods.....	68
Study area.....	68
Statistical approach .....	72
Results.....	75
Livestock and fire .....	76
Vegetation and fire.....	76
Climate and fire.....	77
Relationships among spring vegetation, climate, and livestock .....	78
Local and broad-scale climate .....	78
Multiple regressions .....	79
Structural equation modeling .....	79
Discussion .....	80
Livestock.....	80
Vegetation .....	81
Climate .....	82
Acknowledgements.....	85
Bibliography .....	85

Chapter 3: Rapid vegetation change after land use changes and increasing wildfire activity in arid ecosystems of Southern Russia.....	109
---	-----

Abstract .....	109
Introduction.....	111
Methods.....	114
Study area.....	114
Study period .....	116
Data.....	116
Analysis.....	119
Results.....	121
Greenness and landcover types .....	121
Landcover changes.....	121
Association with permanent sands and burning.....	122
Discussion .....	123
Conclusions.....	127
Acknowledgements.....	128
Bibliography .....	129
Chapter 4: Saiga habitat selection and habitat distribution in southern Russia .....	152
Abstract .....	152
Introduction.....	154
Methods.....	157
Study area.....	157
Data.....	158
Habitat suitability modelling.....	160
Results.....	161

Overall variable importance.....	162
Overall habitat distribution .....	162
Relationships between variables and habitat suitability in the overall model .....	163
Annual and seasonal variability in variable importance .....	163
Annual and seasonal variability in habitat distribution.....	164
Discussion .....	164
Acknowledgements.....	168
Bibliography .....	170

## List of figures:

Figure 1 Total sheep population in three administrative regions of the Republic of Kalmykia (Chernozemelsky, Iki-Burulsky, Lagansky) which closely correspond to the boundaries of our study area (ROSSTAT 2003, 2007).....	53
Figure 2. Study area location (hatched polygon with black outline, grey line – Republic of Kalmykia; the grayscale background represents shaded relief (SRTM30). ....	54
Figure 3. Feathergrass ( <i>Stipa</i> sp.) (top) and sage ( <i>Artemisia</i> sp.) (bottom) dominated communities in the beginning (left) and the end (right) of the vegetation season.....	55
Figure 4. 16-day MODIS reflectance values in bands 1 (triangles) and 2 (squares) for a sample area (median of 9 pixels) affected by fire (hatched line denotes the fire date). Winter-time snow contaminated values are not shown. ....	56
Figure 5. Estimated burned area according to the AVHRR analysis (solid line), validation dataset (dotted line, triangles) and official burned area data (dashed line, squares).....	57
Figure 6. An example of AVHRR-based burned area map for year 2000 (red polygons) comparing with MODIS-derived burned area validation data (outlined in black) shown on top of the AVHRR image from Aug 1, 2000. The boundary of analysis region is shown in magenta.	58
Figure 7. Example of before-burning and after-burning in a Landsat TM image of September, 1988 and an ETM+ image of July, 2001 (5-4-3 band combination). Ubiquitous fire-scars bounded by roads in the 2001 image, most of these scars are from previous years (marked with arrows). The photo represents an area burned in September, 4, 2005, photo taken March, 18, 2006.....	59
Figure 8. Number (line) and area (bars) of burn areas from 2000 to 2007 summarized by size class.....	60

Figure 9. Fire seasonality, i.e., the burned area by month, based on the MODIS image interpretation. ....	61
Figure 10. Burned area mapped for selected years with high fire activity. Gray shading – AVHRR-based burned area maps, black outlines – validation burned area maps (1998 – RESURS-based, others – MODIS-based).....	62
Figure 11. Sheep population and burned area trends in the study area.....	102
Figure 12. Study area boundary and location. ....	103
Figure 13. Relationships between selected explanatory variables and log-transformed burned area . Solid dots and solid line represent values and regression line for period ‘before’ (1986-1996), hollow circles and dashed line represent values for period ‘after’ (1997-2006). ....	104
Figure 14. Change in (a) burned area, (b) maximum August temperatures, (c) livestock numbers, and (d) April NDVI between ‘before’ and ‘after’ periods.....	105
Figure 15. Correlation fields of broad-scale climatic oscillation patterns and local climate during fall months. Red indicates positive correlations, blue negative correlation, light colors - no statistical significance, and dark colors - statistical significance at 0.05 level, black – not analyzed. ....	106
Figure 16. On top, (a) the final structural equation model, and below (b) full model with pathways and variables that were not retained (shown as dashed lines). Straight single-headed arrows represent significant effects of one variable on another ( $\alpha = 0.05$ ), while curved double-headed arrows represent correlations between variables. The relative strength of the effect is indicated by a path coefficient. Only pathways that are significant at $\alpha = 0.05$ are included. See Table 1 for notations and abbreviations.....	107



Figure 17. Average annual decline in total sheep from 1992 to 2008 for the ten countries with the strongest absolute declines. Number in brackets indicates time span in years between the lowest and the highest sheep number during this period (FAO 2010). .....	142
Figure 18. Study area location (hatched area with black outline) in southern Russia .....	143
Figure 19. Trends in spring NDVI and precipitation (sum of March-June period rainfall, mm) in the study area from 1982 to 2007 period. While there were strong increases in NDVI, there was no systematic trend in precipitation. ....	144
Figure 20. Examples of different vegetation types, from top left clockwise: <i>Stipa capillata</i> dominated community, <i>Artemisia lericheana</i> dominated community, community dominated by annual <i>Bromus tectorum</i> and perennial <i>Stipa sareptana</i> , permanent bare sands. Images taken by M. Dubinin in late spring. ....	145
Figure 21. Phenological differences for the three pure vegetation types based on field data and MOD13A1 NDVI in 2007, error bars indicate 1 standard deviation ( $N_{\text{perennials}} = 6$ , $N_{\text{shrubs}} = 6$ , $N_{\text{annuals}} = 6$ ). ....	146
Figure 22. Differences in brightness (0-255, Landsat 5/TM May 5, 2007) for different vegetation communities, $N_{\text{shrubs}} = 51$ , $N_{\text{annuals}} = 26$ , $N_{\text{perennials}} = 98$ . ....	147
Figure 23. Total area in different land cover classes in 1985, 1998, and 2007. ....	148
Figure 24. Association between former permanent sands (red polygons) and landcover (green – grasslands, grey – shrublands, yellow – permanent sands, blue - water) in late spring of 1998 (left, top) and late spring of 2007 (left, bottom). Also shown fragments of Landsat TM images, RGB combination: bands 5, 4, and 3 (right) .....	149

Figure 25. Annual NDVI trends for the years of selected satellite images in a) 1985, Landsat images recorded on May 8 <sup>th</sup> and June 6 <sup>th</sup> and 1998 April 26 <sup>th</sup> and June 13 <sup>th</sup> , and b) 2007 (March 10 <sup>th</sup> and May 5 <sup>th</sup> ).	150
Figure 26. Land cover classifications for 1985 (top left), 1998 (top right), and 2007 (bottom right). White areas were masked out because of cloud contamination in at least one of the images. The apparent striping is caused by Landsat problems with the scan line correction procedures which caused stripes of missing data.	151
Figure 27. Dynamics of livestock and Saiga antelope populations from 1952 to 2008 in Kalmykia.	182
Figure 28. Study area location in southern Russia. Dashed lines in the study area represent protected areas (left - Stepnoi Nature Preserve and right - Chernye Zemli Nature Reserve), hatched area with black outline – study area, thick grey line – administrative boundary of Republic of Kalmykia.	183
Figure 29. Distribution of selected variables. Clockwise from the top left: elevation (meters), burning frequency (number of times a location burned), distance to water sources (thousands of meters), distance to farms (thousands of meters), vegetation communities in 2007 (for legend see Methods section), mean total NDVI (index value). Dashed lines represent boundaries of protected areas.	184
Figure 30. Habitat suitability map using overall dataset showing the point-wise mean. Blue represents preferred habitat, brown non-preferred, white missing data. Violet dots represent Saiga antelope locations used as training points, green - locations used for validation. Dots represent one of the replications and provided as an example. Dashed lines represent boundaries of protected areas.	185

Figure 31. Annual distribution of suitable habitat, averages over 5 replications are shown.	186
Figure 32. Difference of the habitat suitability index for a given year minus the habitat suitability index for all years (2003 – 2007). .....	187
Figure 33. Seasonal distribution of suitable habitat, averages over 5 replications are shown.	188
Figure 34. Response curves showing relationship between the particular variable and logistic output (probability of presence) according to the global model. Gray lines indicate minimum and maximum from 5 replications and thick black line represents a mean. See Table 1 for explanations of units of x-axis. ....	189

## List of Tables:

Table 1. Annual burned area estimation and validation results for study period. ....	51
Table 2. Detection rates of burned areas in different size class categories.....	52
Table 3. Pearson correlations between explanatory variables and burned area for the ‘before’ (1983-1996), and ‘after’ period (1997-2006), and the entire time series, and the stability (i.e., the mean and standard deviation) of the correlation coefficients when moving the split year (1992-1997). T-test shows p-value for significance of the differences between before and after datasets. Significance levels of the univariate correlations between the explanatory variables and burned area are denoted as: ** < 0.01, * < 0.05. Notation and abbreviations: °- variable lagged one year back, WR – West Russia, AO – Arctic Oscillation, NAO – North Atlantic Oscillation, tmx – maximum temperature, pre – precipitation, NDVI – Normalized Difference Vegetation Index. Numbers indicate period or specific month. ....	93
Table 4. Five best multivariate models explaining variability in log-transformed burned area a) with b) without broad climate variables (r denotes Pearson correlation between variable 1 and variable 2, significance levels: ** < 0.01, * < 0.05). ....	95
Table 5. Correlations between fire, local climate measures and oscillation indices AO, NAO, EAWR (monthly values, summarized by season: W – winter, Sp – spring, Su – summer, F - fall and yearly average), number indicates month (significance levels: ** < 0.01, * < 0.05). ....	96
Table 6. Relationships between a) April and b) May NDVI and other variables.....	98
Table 7. Statistical significance of difference in slopes. Dummy – statistical significance for keeping 0/1 variable representing periods ‘before’ and ‘after’. Slopes – statistical significance for different slopes. Significance levels: ** < 0.01, * < 0.05.....	100

Table 8. Results of two-variable best-subset analysis; only variables which participated in more than 1 model are shown. ....	101
Table 9. The definition of nine vegetation associations and other land cover classes for each year based on the classification of four land cover classes for corresponding early and a late spring classifications. ....	137
Table 10. The association of different land cover classes in 2007 with the cumulative burned areas (1998 – 2007). Subtotal for perennial grasslands is the sum of the three classes that had perennials (annuals-perennials, shrubs-perennials, perennials), the subtotal for shrubs is the sum of two shrub classes (annuals-shrubs, shrubs). ....	138
Table 11. The proportion of each land cover class that burned once, twice, etc. up to seven time. ....	139
Table 12. Land cover transitions for two periods: a) 1985 – 1998, b) 1998 – 2007, stable areas are shown in bold. Reported as the area (km <sup>2</sup> ) and the percentage in brackets. ....	140
Table 13. Predictors used to model suitable habitat and determine its drivers and variability in Southern Russia. ....	175
Table 14. Amount of data used for model construction and resulting performance of the models. SD AUC represents average standard deviation across 5 replications, while each SD AUC is calculated by folding and using remaining folds for cross-validation, repeating for every replication. ....	176
Table 15. Variable contribution measured as permutation importance, percentages, averaged over 5 runs: a) overall and annual models, b) seasonal models ....	177
Table 16. Results of jackknife test for variable importance as AUC on test data sorted by sum, as percentages, averaged over 5 runs: a) overall and annual models, b) seasonal models .....	179

Table 17. Amount of suitable and highly suitable habitat and its change across years and seasons in percents of total study area. ....	181
---	-----

## Introduction

Human land use is widely acknowledged to be the most influential disturbance agent (Foley et al. 2005, MEA 2005). Changes in disturbance regimes profoundly affect components of natural and coupled socio-ecological systems such as land cover, ecosystem processes, and biodiversity (Turner 2010). In upcoming decades, land use will likely continue to be more important than other factors of change such as climate (Sala et al. 2000). This is worrisome, because excessive amounts of disturbance may lead to catastrophic shifts in the composition and function of communities, as their critical thresholds of disturbance are reached (Scheffer et al. 2001).

Human land use is both a cause and effect of current and past socio-economic conditions. Massive institutional changes can lead to changes in social and economic conditions of similar magnitude, and as a result, human land use may be altered as well. While often having degrading effects on the environment, massive institutional changes also represent a promising research opportunity, because they can provide natural experiments at an unprecedented scale (Diamond 2001).

After the breakdown of the USSR, land use in the arid grasslands of southern Russia changed substantially, the major change being a strong decrease in livestock numbers (FAO 2010). Like other natural systems, arid ecosystems are quite susceptible to human land use. While rarely used for row crops, arid ecosystems often suffer from degradation due to overgrazing that may lead to significant changes in vegetation and fauna, and can even result in desertification and famine (van de Koppel et al. 1997). The rapid rates of livestock decline in southern Russia in the 1990s were globally unprecedented, and matched by few other places in

the world, including Kazakhstan (FAO 2010), which underwent the same institutional change. However, despite the massive extents and rapid rates of land use changes in the arid grasslands of southern Russia, the environmental consequences and effects of these changes have barely been studied, especially at broad spatial scales. This is unfortunate, because such declines in land use intensity represent a rare opportunity to understand the relationships among land use, vegetation, wildlife, and other disturbance processes in arid grasslands.

My study area was located in the semi-deserts of Southern European Russia and represented an expansive belt of arid ecoregions extending from Western Russia east through Kazakhstan to Mongolia. Typical vegetation communities representative of the arid grasslands of this region were dominated by feathergrass (*Stipa* sp.) and sagebrush (*Artemisia* sp.). Summers were hot and dry, and winters were cold with little snow. Annual precipitation ranged from 210 to 340 mm and summer droughts were common. About 10% of the study area was protected by the strict nature reserve Chernye Zemli (“Black Lands”). The study area covered about 30,000 km<sup>2</sup> (approx. 2 Landsat scenes).

The overarching goal of my study was to identify the effects of the recent socioeconomic changes on ecosystem patterns and processes, including fire, vegetation and wildlife. The project focused on the area shared by two administrative regions in southern Russia: Republic of Kalmykia and Astrakhan region, where land use changes have been widespread, and where the last European population of saiga antelopes is at risk of extinction. My general research questions were: how, and why have vegetation and fire regimes changed, and what are the relationships among land use, fire, vegetation, climate, and saiga habitat selection. My analysis was based on remotely sensed TM/ETM+, MODIS, and AVHRR satellite data as well as on a variety of field data and thematic maps. The dissertation consists of an introduction, which



provides a summary of the entire dissertation, and four chapters, which examine specific research questions in detail.

## **Chapter 1**

***Research question: How did fire dynamics change before and after the breakdown of the Soviet Union?***

***Hypothesis: Fire occurrence increased significantly after the collapse of the Soviet Union.***

Fire is a prominent disturbance factor in arid grasslands, but little is known about its history in my study area, or in Central Asia in general. Furthermore, no prior studies have examined how fire dynamics change after a massive socio-economic shock. Hence the collapse of the Soviet Union may provide a case study that is relevant for arid grasslands globally.

To reconstruct burned area I used AVHRR data acquired from NOAA's Comprehensive Large Array-data Stewardship System (CLASS) for the period of 1985-2007 covering April until August annually. From the full set of images, two were selected for every year, with one representing the beginning and the other representing the end of the active burning season (late April – late August). Every image had low cloud cover and was free from geometrical and radiometric errors. Each image pair was combined in annual multiband composite, with four bands from the 'before' image and four bands from the 'after' image. For every year, burned and non-burned areas were identified using visual photo-interpretation based on specific spectral responses and visible changes in two images. All samples were combined into one comprehensive training dataset.

We performed image classification using decision trees. We used training data to parameterize the classification and the resulting decision trees were used to estimate burn probability for each pixel. Decision trees are a nonparametric method, i.e., no assumptions of underlying distribution of the data were made. To generate more stable models, we used bagging approach to derive an ensemble of tree classifiers. We trained each tree classifier separately using a 50% random subset from the comprehensive training dataset. The set of 30 trees was used to classify image stacks for every year.

Results provided strong evidence that burned area increased significantly in recent years compared to the beginning of the series in 1985. I observed a dramatic change in burned area starting in 1997-1998 and burning has remained widespread ever since. Since 1997, burning exhibited significant variation among years compared to the period before 1997.

We conducted validation for 13 out of the 23 years, using separate burned area datasets, generated from various remote sensing datasets, such as Landsat/TM, ETM+ (1986-1989 and 1999), RESURS/MSU-SK (1996-1998), TERRA/MODIS (2000-2007). Overall, the AVHRR burned area map was about 60-70% accurate, but showed some underestimation of burned area, because of low-resolution bias of the AVHRR data. However, established temporal trends were robust, and matched official data well, highlighting even more the conservative nature of governmental information. We conclude that institutional change can lead to significant changes in burning in arid grasslands and these changes that are lagged in time.

**Related paper** (published): Dubinin, M., Potapov, P., Lushchekina, A., & Radeloff, V.C. (2010). Reconstructing long time series of burned areas in arid grasslands of southern Russia by satellite remote sensing. *Remote Sensing of Environment*, 114, 1638-1648

**Related paper** (published): Dubinin, M., Lushchekina, A.A., & Radeloff, V.C. (2010).

Assessment of modern burning regime in arid ecosystems using remote sensing data (case study of Chernye Zemli) [in Russian]. Aridnye Ecosystemy

## Chapter 2

**Research question:** *How is the annual burned area related to changes in livestock, vegetation and climate?*

*Hypotheses:* The burned area in a given year was:

- 1) positively correlated with the amount of vegetation during the spring growing season immediately prior to the summer fire season, because of higher fuel availability;*
- 2) positively correlated with the amount of vegetation during the secondary growing season in the fall of the previous year, again due to higher fuel availability;*
- 3) positively correlated with climatic conditions that are favorable for vegetation growth during spring and fall of the previous year, also due to higher fuel availability;*
- 4) significantly more correlated with climate conditions during the period after livestock numbers had declined and grazing pressure lessened, than during the period before.*

Our results from the first chapter highlighted changes in the burning regime that were linked to institutional change and the resulting land use changes. However, the actual environmental drivers of this change were not quantified. Though we knew that livestock numbers collapsed and vegetation started to recover, we did not know if land use or climate change was the main reason for the increase in fire activity, and how all these factors were

interrelated. The main goal of my second chapter was thus to identify the interactions among these drivers and their effects on burning dynamics.

To address the research question, we conducted several statistical analyses of the significance of the relationships among annual burned area and different human and environmental variables. Burned area estimates from Chapter 1 were correlated with a number of environmental variables, such as local precipitation, maximum and average temperature, and broad-scale climate indices (West Russia, Arctic and North Atlantic Oscillation indices), vegetation (represented by GIMMS NDVI for April, May and August aggregated over study area and their previous year values), as well as the main human driver, represented by livestock presence. To analyze the relationship of my burned area estimates with these different environmental variables before and after institutional collapse, the dataset was separated into two periods (before and after) and correlations were calculated for each period separately. This analysis was also coupled with regression modeling to try to assess if the slopes of the relationships were different before and after fires had started. Finally, derived relationships were used to build a structural equation model that enabled the analysis of the whole set of relationships simultaneously, and removed weakly related variables.

When analyzed over the entire time period, burned area was significantly negatively correlated with livestock numbers and positively correlated with aggregated NDVI value in April (spring greenness). The highest positive correlation of burned area was with NDVI in May, meaning that more biomass during this month resulted in more fires. Moderate correlations were found with climatic variables such as precipitation in April and maximum August temperatures, highlighting the presence of direct and indirect effect of these factors on vegetation and fire.

When analyzing the two time periods separately, amount of livestock was not an important factor. Consistent with the overall analysis, aggregated NDVI values in April were highly correlated with burned areas both before and after the collapse. These results provided an interesting quantitative picture of the dynamics of the ecosystems and the relationships of its components with burned areas. The main determinant of burning was May NDVI (right before the burning season starts). Interestingly, although April NDVI was also significantly positively correlated with burning, the direct link with it was not significant, illustrating the fact that a) April pastures are heavily grazed, and b) vegetation in May is dependent of vegetation in April. None of the precipitation measures were directly significant in explaining burning, and this suggests that moisture was not important, probably because of precipitation during the fire season in the summer. On the other hand, precipitation was not eliminated from the structural equation models, and linked to vegetation in April and May.

With the exception of fall greenness, our results could not reject any of our null-hypotheses. Livestock represented an important background disturbance factor, but was not significant explanatory factor if two sub-periods were concerned separately. Interestingly, there was a very substantial increase in the correlation with May NDVI in the period after the collapse. This strong increase in correlation May NDVI suggested that because livestock removed most of the vegetation, NDVI was not a good predictor of burning before the collapse, but NDVI became an important predictor after the collapse as vegetation recovered.

Our results highlight a complex system of interacting drivers that can jointly explain the burning regime. Socio-economic changes unmasked the effect of climate on burning after the collapse, while during Soviet times, fires were heavily suppressed and fuel loads were low. Our analysis showed that the broad climate oscillation indices that are often used for explanation of

burning regime should be used with care, as it they are not always related to local climate and thus the mechanisms of their effect on burning were unclear.

**Related paper** (in review): Dubinin, M., Lushchekina, A., & Radeloff, V.C. (2010).

Relationships between climate, livestock, and fire in the arid grasslands of southern Russia.

Ecosystems.

## Chapter 3

***Research question: What are the effects of fire and past land-use on vegetation communities in semi-deserts?***

*Hypotheses:*

- 1) the area dominated by grasses increased and the area dominated by shrubs declined;*
- 2) the area of bare, open sands decreased; and*
- 3) areas where grasses replaced shrubs were associated with both wildfires and formerly open sands.*

In Chapter 1 and Chapter 2, we found significant increases in fires and this increase was related to vegetation. However, we used only a proxy variable for vegetation (NDVI), and I did not know if the increases in fires resulted in – or were caused by – changes in the composition of the vegetation community. The goal of this chapter was to understand the effects of land use change on vegetation communities in semi-deserts.

Vegetation changes can reflect both recent changes and long-term land use legacies. The data on the extents and dynamics of disturbance regimes (fire and grazing) makes it possible to relate changes in the disturbance regime changes in to vegetation composition. The underlying question was whether the effect of these disturbances was substantial enough to result in changes in vegetation over a large area. However, the resulting vegetation patterns may also be affected by past land-use legacies, such as unsustainable row crops production attempts that resulted in rapid wind erosion and desertification. In this chapter we thus asked foremost what kind of vegetation community changes have occurred, and secondly if past land-use signals and fires were more common in areas now occupied by perennial grasslands.

To conduct the vegetation classifications we exploited the specific phenological patterns of the different vegetation communities. Annual grasses had more pronounced peak in greenness (NDVI) in early spring (March-April), perennial grasses in late spring (May), and sagebrush had a weaker NDVI signal overall, but its growing season was longer. We used these phenological features and carefully selected Landsat scenes (one for early spring and one for late spring for every year) to conduct classifications of the dominant vegetation types for 1985, 1998, and 2007.

The mapping of the burned areas in Chapter 1 allowed us to examine how fires affected vegetation patterns. Classification results showed considerable increase in perennial grass-dominated communities at the expense of shrub-dominated in the study area from 1985 to 2007. Conversion took place in almost 20% of the study area and more than 60% of these changes happened within boundaries of burned areas. However, a substantial portion of the study area showed an increase in perennial grass-dominated communities even without burning. Also, areas of change were related to formerly plowed fields, but not limited to them.

The increase in grasslands suggested a positive feedback between perennial grasses and fire. Areas with more fires were more likely to convert to grasslands, and grasslands were more likely to burn. Our results also supported conclusions of plot-level studies of vegetation succession conducted by the local scientists who reported that a type conversion from shrubs to grasses can also be caused solely by release of grazing pressure. However, we found that the change was greatly facilitated by fire, and areas that had burned more than three times since 1998 were more likely to be covered by perennial grassland vegetation compared to areas that burned less frequently.



**Related paper** (to be submitted): Dubinin, M., Lushchekina, A., & Radeloff, V.C. (2010). Rapid vegetation change after land use changes and increasing wildfire activity in Southern Russian semi-deserts.

## Chapter 4

**Research question:** *What constitutes suitable habitat for Saiga antelopes in our study area and how does it change among years and seasons?*

*Hypotheses:*

- 1) there is no significant difference among years in terms of the amount and the patterns of suitable habitat;*
- 2) there are substantial differences in preferred habitat among seasons in terms of the amount and the patterns of suitable habitat; and*
- 3) vegetation type and productivity are among of the most important driving factors of habitat selection by Saiga.*

The Saiga antelope is the last free roaming antelope of Europe, and the last wild population in Russia is located in my study area, and has declined precipitously in recent decades. The sharp decline of Saiga populations was usually attributed to poaching, but landscape change may also play an important role. However, effects of increases in fires and vegetation change affected Saiga habitat are unclear. For my fourth chapter, my goal was to describe habitat selection by Saiga especially in relation to the environmental changes that I reported in my second (fire) and third (vegetation) chapters.

My fourth chapter examines habitat selection at a local scale. The analysis was limited to a subset of the study area from the previous chapters which overlapped with two protected areas in the Republic of Kalmykia and in Astrakhan region. I used maximum entropy modeling and a used-versus-background approach to estimate probabilities of saiga presence. Saiga occurrence data was collected from 2003 to 2007 by the staff of the reserves and used together with maps of various human and environmental variables to estimate what variables best explain the variability in the occurrence data, and how explanatory power varies among years and seasons. Among other variables, fire perimeters and fire frequency from Chapter 1, as well as the vegetation classifications from Chapter 3 were included.

I found a strong relationship between fires and habitat selection by Saiga. Areas that burned more often were consistently selected more frequently by saiga. Another very important factor that contributed substantially to most of the models was the distance to water sources. On the other hand, the effects of vegetation structure and composition were not particularly strong in my study, which is inconsistent with local field-based studies on saiga diet. Areas close to the farms were also avoided, indicating competition with livestock.

Our results highlight that changes in fire dynamics likely affected habitat selection by Saiga antelope. The clustering of Saiga in fire affected areas is probably an indication that poaching by humans and predation by wolves are still significant issues, even in the areas that are protected, so that Saiga antelopes select areas with high visibility in order to avoid threats. This information is new and has implications for management of this endangered species.

**Related paper** (in preparation): Dubinin, M., Lushchekina, A., & Radeloff, V.C. (2011). Habitat selection by Saiga antelope in Southern Russia and its inter- and intra-annual variability.

## Significance

The results of my research add to the understanding of the relationships between land use change, fire, vegetation and wildlife dynamics after dramatic institutional changes. These results contribute to science and conservation in three ways: they provide new ecological insight, they inform management and conservation recommendations, and they make methodological advances.

**Ecologically**, my work addresses the general problem of coupled dynamics of different environmental factors, both human and natural as well as changes in the disturbance regimes. My study is of particular interest because it deals with an environment that underwent a substantial decrease in land use intensity and the environment is currently recovering, which is rather rare, since land use is generally intensifying globally. My results help to understand historical and modern dynamics of fire, vegetation and grazing and provide statistical evidence for the drivers of these patterns in arid ecosystem. From the applied point of view, my research provides a foundation to study habitat selection of Saiga antelopes, which was not possible before, as neither fire nor vegetation effects were estimated explicitly over large enough areas in the region of study. More specifically my study:

- *Provides novel information about a unique ecosystem and its species.* My research fills an important information gap about arid ecosystem and species of southern Russia, some aspects of which were not represented in scientific literature before. Comparable ecosystems exist in other part of the world (for example, the mixed grasslands of the

Great Plains, USA), but they have very different disturbance dynamics due to a much higher level of human domination.

- *Provides evidence of substantial and widespread environmental changes occurring at a regional scale.* Though my study area was small, it is representative for the vast arid belt which traverses the countries of Central Asia and Mongolia, all of which underwent similar socio-economic changes as Kalmykia. My research provides the first evidence of both an increase in fire activity and vegetation change following the collapse of the Soviet Union. Preliminary evidence suggests that processes of change in burning regimes and vegetation were similar in the entire region.
- *Provides evidence for the existence of a restorative threshold effect.* Changes in ecosystem processes, especially fire, were rather abrupt, suggesting the presence of a tipping point, after which rare events increased in frequency and in magnitude. Further monitoring of fires in the future could provide even more information about this basic ecological mechanism. Burned area might serve as an important indicator of coming future changes if increasing livestock numbers reach the tipping point again, causing the ecosystem to revert back into the pre-fire state. A possible indication of that was the year 2010, which was exceptionally dry, and exhibited an unprecedented amount of burning throughout European Russia, but **not** in my study area, potentially because livestock numbers have increased again.
- *Highlights interactions among disturbances.* My research highlights the importance of interactions among fire, climate, and livestock, as well as land use legacies. My results show that gradual changes in vegetation introduced by the decrease of one disturbance

agent (livestock) can be amplified by a switch to another disturbance agent, and how vegetation interacts with particular past land use.

From the **management and conservation** point of view, ecosystem and species conservation can benefit from a better understanding of ecosystem structure and dynamics, and species habitat selection that my research provides. Some results of my project will have direct management implications. For example, I suggest establishing additional artificial water sources for saiga, to better control poaching, and to experiment with additional livestock grazing in the protected areas.

My research also highlights the need for better monitoring programs for fire, Saiga antelopes and vegetation to quantify the ecological patterns and processes. Better data could also improve the models that I have created and further improve understanding. Hopefully, my research will inform the development of conservation strategies aimed at Saiga antelope currently underway in the region.

Last but not least, some of the technical approaches introduced by this research represent **methodological advances** and provide new avenues to study changes in land cover and burning using remote sensing in semi-desert ecosystems. My results highlight how machine-learning approaches such as decision trees, support-vector machines, and maximum entropy can make important contributions to both remote sensing analyses and wildlife habitat modeling in arid grasslands. The use of decision trees is promising for the analysis of long time series of remote sensing data. Common set of models applied uniformly to the sequence of images can help to reconstruct long-term times series in more robust way, especially when dealing with data that is

heterogeneous in quality. A phenological approach to image selection for classification provides effective methods for discrimination of vegetation associations, otherwise hard to distinguish, like grasses and shrubs. Structural equation modeling provides more comprehensive view of the ecosystem, because it integrates pair-wise correlations between sets of variables in an overarching framework.

## **Bibliography**

Diamond, J. 2001. Ecology: Dammed Experiments! *Science* 294:1847-1848.

FAO. 2010. FAOSTAT, <http://faostat.fao.org>

Foley, J. A., R. DeFries, G. P. Asner, C. Barford, G. Bonan, S. R. Carpenter, F. S. Chapin, M. T.

Coe, G. C. Daily, H. K. Gibbs, J. H. Helkowski, T. Holloway, E. A. Howard, C. J.

Kucharik, C. Monfreda, J. A. Patz, I. C. Prentice, N. Ramankutty, and P. K. Snyder. 2005.

Global consequences of land use. *Science* 309:570-574.

MEA. 2005. Millennium Ecosystem Assessment. Global Assessment Reports. Volume 1:

Current State & Trends. Island Press, Washington, DC.

Sala, O. E., F. S. Chapin, III, J. J. Armesto, E. Berlow, J. Bloomfield, R. Dirzo, E. Huber-

Sanwald, L. F. Huenneke, R. B. Jackson, A. Kinzig, R. Leemans, D. M. Lodge, H. A.

Mooney, M. Oesterheld, iacute, N. L. Poff, M. T. Sykes, B. H. Walker, M. Walker, and D.

H. Wall. 2000. Global Biodiversity Scenarios for the Year 2100. Pages 1770-1774 *in*.

Scheffer, M., S. Carpenter, J. A. Foley, C. Folke, and B. Walker. 2001. Catastrophic shifts in ecosystems. *Nature* 413:591-596.

Turner, M. G. 2010. Disturbance and landscape dynamics in a changing world. *Ecology* 91:2833-2849.

van de Koppel, J., M. Rietkerk, and F. J. Weissing. 1997. Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends in Ecology & Evolution* 12:352-356.

# **Chapter 1: Reconstructing long time series of burned areas in arid ecosystems of Southern Russia by satellite remote sensing**

## ***Abstract***

Fire is an important natural disturbance process in many ecosystems, but humans can irrevocably change natural fire regimes. Quantifying long-term change in fire regimes is important to understand the drivers of changes in fire dynamics, and the implications of change in fire regime for ecosystem ecology. However, assessing fire regime changes is challenging, especially in grasslands because of high intra- and inter-annual variation of the vegetation and temporally sparse satellite data in many regions of the world. The breakdown of the Soviet Union in 1991 caused substantial socioeconomic changes and a decrease in grazing pressure in Russia's arid grasslands, but how this affected grassland fires is unknown. Our research goal was to assess annual burned area in the grasslands of southern Russia before and after the breakdown. Our study area covered 19,000 km<sup>2</sup> in the Republic of Kalmykia in southern Russia in the arid grasslands of the Caspian plains. We estimated annual burned area from 1985 to 2007 by classifying AVHRR data using a decision tree algorithm, and validated the results with RESURS, Landsat and MODIS data. Our results showed a substantial increase in burned area, from almost none in the 1980s to more than 20% of the total study area burned in both 2006 and 2007. Burned area started to increase around 1998 and has continued to increase, albeit with high fluctuations among years. We suggest that it took several years after livestock numbers



decreased in the early 1990s for vegetation to recover, to build up enough fuel, and to reach a threshold of connectivity that could sustain large fires. Our detection algorithm for burned areas was effective, and captured burned even areas with incomplete annual AVHRR data. Validation results showed 68% producer's and 56% user's accuracy. Lack of frequent AVHRR data is a common problem and our burned area detection approach may also be suitable in other parts of the world with comparable ecosystems and similar AVHRR data limitations. In our case, AVHRR data were the only satellite imagery available with time series long enough to reveal marked increases in fire regimes in southern Russia before and after the breakdown of the Soviet Union.

## ***Introduction***

Fire is one of the main disturbance agents in grasslands and savannahs. Fire shapes vegetation structure and composition, and represents an important land-management tool (Pyne 1984). Grasslands, woody savannahs, and savannahs represent more than 60% of the global burned area (Tansey et al. 2004), and in regions with high aridity such as Central Asia, grassland fires account for 80% of all active fire counts (Csiszar et al. 2005). Fuel loads and emissions from grassland burning are relatively small (van der Werf et al. 2006), but grassland fires can foster the spread of invasive species (Brooks et al. 2004), affect wildlife (Archibald and Bond 2004), and cause air pollution that can spread as far as the Arctic (Stohl et al. 2007). Furthermore, interactions between grassland fires and human land use may result in ecosystem degradation, hydrologic changes, soil disturbance, and shrub encroachment (Archer et al. 1995). The restoration, conservation, and management of arid grasslands thus require solid information on their fire regime and its change over time. Our goal here was to assess fire regime changes in the grasslands of southern Russia before and after the breakdown of the Soviet Union.

Fire regimes in general are largely determined by fuel moisture, fuel amount, and ignition sources (Bond and van Wilgen 1996). Arid grasslands are biomass-poor, and at least seasonally dry ecosystems, in which fuel amount and connectivity are the main limiting factors for fires (Meyn et al. 2007). Climate change can cause long-term changes in fuel amounts, but land use affects fuel amounts more directly and acutely. The main control of fuel amount in many arid grasslands is livestock (Bahre 1991). Unlike mesic grasslands, which evolved with intensive mammalian herbivory, arid grasslands are more sensitive to livestock grazing (Mack and Thompson 1982) often leading to overexploitation and degradation (Akiyama and Kawamura 2007; Zhang et al. 2007). Vegetation can recover when livestock grazing decreases, and the resulting increase in

fuel amounts and connectivity may be able to sustain extensive wildfires (Liedloff et al. 2001). However, fire increases resulting from a decrease in livestock grazing have rarely been studied, and the long-term relationship between livestock grazing, burning dynamics, and environmental factors is poorly understood. Thus, it is often inappropriate to study livestock-fire relationships with small plots and difficult to employ a classical experimental design at such broad temporal and spatial scales.

Where controlled experiments are impossible, natural experiments may offer insights (Diamond 2001). Significant socioeconomic disruptions and the resulting land-use changes can provide natural experiments to study accompanying environmental changes (Kuemmerle et al. 2007). Research on land-use change typically examines how socioeconomics affect land use directly (Lambin and Geist 2006). However, indirect cascading effects, such as alterations of disturbance regimes, may have a much larger effect on the ecosystem (Burcher et al. 2007). In the context of livestock grazing and grassland fires, the ideal natural experiment would comprise a strong decrease in livestock grazing, sufficient time for the resulting changes in fire regimes to manifest themselves, and relatively constant conditions for all other relevant factors. The breakdown of the Soviet Union in 1991 provided such a natural experiment.

The former Soviet Union including Russia experienced dramatic changes in livestock populations in the arid grasslands, such as in the Republic of Kalmykia (Figure 1). In the early 1970s, under state-controlled, plan-based economic policies, sheep numbers more than doubled, and stayed at over 800 thousand head for about twenty years. During these years, the bulk of sheep industry was represented by large agricultural enterprises, which overexploited the grasslands. Grazing intensified up to a point where some pastures were grazed year-round. Intensive grazing caused widespread wind erosion and vegetation degradation (Zonn 1995). Due to overgrazing, southern Russia was called “Europe’s first anthropogenic desert” in the mid 1990s

(Saiko and Zonn 1997). Some model forecasts predicted that even with protection, more than 66% of southern Russia's grassland would convert to bare sand by 2000 (Vinogradov 1995), and substantial governmental action was taken to combat desertification.

However, following the breakdown of the USSR in 1991, livestock populations dropped by almost an order of magnitude (Figure 1) and remained low until around 2000 (Brooks and Gardner 2004; ROSSTAT 2007). Large collective and state-owned farms were no longer subsidized after 1991, resulting in broad-scale de-collectivization and abandonment. The livestock declines allowed vegetation to recover and likely caused an increase of grassy fuels. Thus the hypothesis is that the decrease in grazing pressure resulted in an increase in grassy fuels, and ultimately in an increase in grassland fires. The problem is that no fire data are available to test this hypothesis since official fire records for the region, similar to other regions are incomplete and inaccurate (Soja et al. 2004).

Remote sensing and satellite imagery have been used for more than 20 years to monitor fires in many different parts of the world (Csiszar et al. 2004). Coarse resolution satellite data are one of the best sources for historic burned area mapping because of their high temporal resolution and long history of acquisitions (Arino et al. 2001). Two basic types of algorithms exist to estimate fire areas from Advanced Very High Resolution Radiometer (AVHRR) or Moderate-Resolution Imaging Spectroradiometer (MODIS) data: a) detection of active fires, and b) mapping burn scars or burned areas. Though area burned correlates well with the number of fires, deriving burnt areas from active fire detections is error prone, especially in some ecosystems, because the detected hotspots can underestimate the actual burned area (Giglio et al. 2006; Hawbaker et al. 2008; Miettinen et al. 2007). In contrast, burned area assessments identify postfire disturbance, but not the actual fire events.

Typically burned area mapping involves band differencing and thresholding or classifying single or multitemporal data in the form of raw band values and derivative indices

(Arino et al. 2001; Gong et al. 2006; Kučera et al. 2005). Burned area maps from coarse spatial resolution data are prone to several types of errors: low resolution bias, lack of imagery in certain areas, and geometric and radiometric errors (Barbosa et al. 1999; Boschetti et al. 2004). Most of these problems are more prevalent in AVHRR data, and have been solved or notably reduced in recent sensors such as MODIS on board of Terra and Aqua satellites. However, MODIS images are only available since 2000 and this relatively short time series limits the analysis of long-term trends. Thus AVHRR remains the only remote sensing dataset capable of reconstructing long-term (20+ years) burning trends (Chuvieco et al. 2008; Kučera et al. 2005) at a regional scale. The classification of satellite images for burned area mapping typically requires regular AVHRR observations, composited from daily data. Full resolution AVHRR data (Local Area Coverage, LAC, or High Resolution Picture Transmission, HRPT, both at 1.1 km at nadir) greatly improves the classification of burned areas (Pu et al. 2007; Razafimpanilo et al. 1995; Sukhinin et al. 2004).

Unfortunately, the amount and quality of AVHRR data vary greatly among different regions. Daily AVHRR observations at 1.1 km spatial high resolution are not always available, especially for those parts of the world where local archives had not been established until the late 1990s. Southern Russia is one such area, and only a limited amount of AVHRR data is available in international archives such as NOAA's Comprehensive Large Array-data Stewardship System (CLASS, <http://www.class.noaa.gov/saa/products/welcome>). Because of the limited data availability, existing remote sensing methods designed to map burned areas from AVHRR data can not be applied and there is a need to develop new approaches to map burned areas in regions where only CLASS-type AVHRR data are available.

The main goal of this study was to assess changes in burning in the grasslands of southern Russia before and after the breakdown of the Soviet Union. Our hypothesis was that annual burned area increased substantially after the breakdown of the Soviet Union due to the decline of livestock populations and more abundant fuels. Our second goal was to develop and test a method to derive burned area trends from temporally sparse CLASS-type AVHRR data. Our study was motivated by 1) the absence of any solid information on distribution of fires throughout 1980s and '90s in the study area and throughout Central Asia, 2) substantial socio-economic changes in the region providing a unique 'natural experiment' to examine the effects of decreasing livestock grazing on fire regimes, and 3) the need for a remote sensing approach that can map long-term burned area trends in arid grasslands from sparse AVHRR data.

## ***Methods***

### **Study area**

Our study area was located in the grasslands of Southern European Russia and occupied about 19,000 km<sup>2</sup> of the Republic of Kalmykia and Astrakhan Region (Figure 2). The study area was defined in the west and east by the administrative boundaries of the Republic and in the north and south by the boundaries of the common area of available AVHRR overpasses. The study area was sparsely populated (population density 0.8 to 1.4 persons/km<sup>2</sup> (CIESIN and CIAT 2005) with no expectation for population growth (CIESIN et al. 2005).

The climate of the study area was arid, with hot, dry summers (mean daily temperature of +24°C in July; max +44°C, 280 days of sunshine per year on average). Annual precipitation was 150 to 350 mm (with a mean of 286 mm for 1985 to 2007). Summer droughts were common, and most of the precipitation fell in spring (43% of all precipitation), coinciding with the period of

most vegetation growth (Walter and Box 1983). The topography of the study area is predominantly flat with a mean elevation of -15 m below sea level. The study area had a complex geological history of transgressions and regression of the Caspian Sea and soils characterized by a gradient from sandy aeolian deposits and sandy loams in the southeast corner of the study area to clay loam in northwest corner (Kroonenberg et al. 1997).

Vegetation associations were typical for northern Precaspian Plains and represented combinations of steppe and desert types. The main vegetation associations were shortgrass steppe (*Stipa* spp., *Festuca* spp., *Agropyron* spp., *Anizantha tectorum*, and other graminoids, **Figure 3**) and sage scrub (*Artemisia* spp., *Kochia prostrate*, **Figure 3**) (Golub 1994). Shortgrass steppe was characterized by a short growing season in April and May and rapid senescence in the dry summer. The grasses exhibited fire adaptation due to dense bunches which protect meristems and generate abundant fuels for burning. Sagebrush (*Artemisia* spp.) dominated shrublands have less biomass, but a longer growing season, and sometimes exhibit a second vegetation peak in the fall or even early winter (Kurinova and Belousova 1989). *Artemisia* spp. is more susceptible to fire because its buds are situated above ground and can be killed or damaged by fires. The lack of fire tolerance by *Artemisia* spp. might lead to its substitution by *Stipa* spp. and other graminoids. The primary human land use of the grasslands in the study area was as rangelands to support grazing for domestic livestock, mainly sheep and to a lesser extent cows and goats.

## Study period

We studied burned area dynamics from 1985 to 2007. The choice of the study period was determined by remote sensing data availability and because of significant changes in land use after 1991. The 23 year fire record that we derived was one of longest for a burned area estimation with coarse resolution satellite data. Other burned area studies have either focused on

forests or analyzed smaller time spans while studies of equivalent time span are particularly rare (Chuvieco et al. 2008; Kučera et al. 2005).

## **Data**

### *Classification data*

We used coarse spatial resolution AVHRR level 1B imagery in the form of digital numbers to estimate burned areas and three types of coarse- and medium- resolution satellite imagery for validation. We chose AVHRR for our burned area time series, despite its limited spatial and spectral resolution, because the length of the AVHRR data record matched our research goal to compare pre- and post 1991 burned areas. For the burned area classifications, we analyzed AVHRR LAC images acquired by National Ocean and Atmospheric Administration (NOAA) satellites (9,11,14,16,17). LAC data are stored onboard the satellite and downloaded at one of the NOAA receiving stations. All daytime images from 1985 to 2007 were downloaded from NOAA's CLASS. From these data, we selected a subset of images ranging from April 1<sup>st</sup> to September 20<sup>th</sup> of each year, corresponding to the beginning and the end of the growing season, and fully capturing the time of summer droughts and thus the fire season. From all images for each year, two images were selected according to the following criteria: i) the sensor angle was less than 45° degree off nadir, ii) no clouds obscured the study area, iii) the first image represented the peak of the growing season or close to it (April, May, or early June), and iv) the second image represented the end of the dry season and the end of the fire season (late August to early September). Most of the CLASS AVHRR LAC imagery had to be discarded. Typically, AVHRR images covering whole study area were available for every second day of the study period. For example, in 2006, out of 173-174 days (April-September), 90 were available. Among those 49 images were discarded due to excessive cloud amount, 10 due to scanner malfunction



and 29 due to position of the study area  $45^\circ$  degree off nadir. Among 12 remaining images of good quality, 3 were available for May-Early June and 3 for August. We based our classification on the assumption that fires will raise reflectance, due to the removal of vegetation, and that after fire event brightness will remain elevated through the rest of the season. Persistent increase in brightness was confirmed by preliminary MODIS analyses (Figure 4). Unfortunately, due to limitations in data availability, the dates of the AVHRR images were not identical among years. Image dates for the earlier image with the exception of 1998 (early April) were all from early May or the first week of June. Later image dates ranged from early (1994, 2000, and 2005) to late August and to early September. All satellite images were clipped to the study area boundaries.

The two selected images for each year were then composited into 12 bands, where bands 1-3 represented the AVHRR spectral bands from the spring image (620-670, 725-1100, 3550-3930 nm); 4-6 represented the AVHRR bands from the fall image; 7-9 were band differences where each fall band was subtracted from the corresponding spring band and offset value was added to avoid negative numbers; and 10-12 represented NDVI for the first and second image (NDVI1, NDVI2) and their difference (NDVI12). We removed band 5 as it was not present for all AVHRR instruments. Bands 4 was removed from the classification after preliminary analysis showed that this bands did not help differentiate burned from surrounding, non burned areas. Dark object subtraction was used to correct the images radiometrically (Song et al. 2001). The minimum observed value was selected from the Caspian Sea, and we subtracted the digital number observed over clear water for each band from all pixel values in each respective band.

#### *Validation data*

For validation, we used Terra/Moderate Resolution Imaging Spectroradiometer (MODIS), Landsat/Thematic Mapper (TM), Enhanced Thematic Mapper Plus (ETM+) and RESURS/Multi Spectral Scanner (MSU-SK) imagery. The MODIS Level 1B calibrated radiances product (MOD02QKM) with 250-m resolution was obtained from NASA's Level 1 and Atmosphere Archive and Distribution System (<http://ladsweb.nascom.nasa.gov>) and used for validation from 2000 to 2007. In addition to the validation, we also used MODIS data to estimate the temporal regime of burning (e.g., fire season length), because of MODIS' high temporal resolution. Landsat TM/ETM+ data were obtained from the USGS EROS Data Center at 30-m resolution for 1986 to 1989 and for 1999. No TM/ETM+ data were available in either public (USGS GLOVIS) or commercial archives (Euroimage) for the period of 1990 to 1998. Instead we used RESURS MSU-SK data obtained from R&D center Scanex (<http://scanex.ru>) at 150-m resolution and 4 spectral bands ranging from 540 to 1175 nm for 1996 to 1998. All validation data was collected for the period correspondent to period of the AVHRR data that we used (i.e., April – September). Last but not least, we obtained official fire statistics (Ministry of Emergency). Although the official fire statistics do not include all burned areas, the statistics provided comparison for the recent years of the burned area time series.

## **Data processing**

### *Geometric correction*

For every year in the sequence, the first AVHRR image was co-registered to the MODIS image of June 7<sup>th</sup> 2008 with a second-order polynomial transformation and nearest-neighbor resampling in ERDAS IMAGINE. The second image from a given classification pair was then co-registered to the first image using the same approach. The pixel size was set to 1.1 x 1.1 km. Semiautomatic tie point collection provided an initial set of ground control points (GCP). For

each image from 7 to 10 GCPs were used for registration. Overall co-registration accuracy (root mean square error) was less than one pixel for each image. MODIS and Landsat data georeferencing accuracy is far superior to that of AVHRR and no additional georeferencing was necessary (Justice et al. 1998; Lee et al. 2004). Visual checking confirmed that the georeferencing of the validation data to the AVHRR data and 1 : 200,000 topographic maps was fine. RESURS data were co-registered to the June 7<sup>th</sup> 2008 MODIS image using the same methodology as for the AVHRR data. All imagery was reprojected to WGS84 coordinate system and equal-area Albers projection (central meridian at 45 degrees, first standard parallel 52 and second standard parallel 64 degrees).

### *Training data*

Visual image interpretation provided training data for both burned and un-burned areas for the classification. For each AVHRR image pair, a set of burned and non-burned areas was digitized. Burned areas were identified based on one of two possible features. Areas that had burned late in the season and were still black were characterized by lower reflectance in the fall image, especially in AVHRR band 2 (Pereira 1999). Conversely, areas that had burned earlier in the season were highlight by vegetation removal and a high proportion of bare ground, causing higher reflectance in bands 1 and 2 of the fall image. For every year non-burned areas were also identified, including unburned vegetation, bare ground and water bodies. All samples were combined into one comprehensive training dataset, which consisted of 11,706 pixels (2,629 pixels for burned areas and 8,447 for non-burned). We used a combined training dataset, rather than keeping training data for each year separate, because we could not detected any visible burned areas in the AVHRR images in the 1980s.

### *Classification*

Our classification method had to account for the fact that there were major limitations in the AVHRR image availability (see above). We used decision tree classification to calculate the certainty that a given pixel burned in a given year. Decision trees are a robust approach to classify satellite data (Friedl and Brodley 1997; Hansen et al. 2000) and are effective tool for burned area mapping in grasslands (Maggi and Stroppiana 2002; Stroppiana et al. 2003). Decision trees are nonparametric, i.e. no assumptions about the underlying distribution of the data are made, and decision trees can capture non-linear relationships between spectral data and different information classes (Breiman 1984). To create more generalized and stable classification models we used a bagging approach (Breiman 1996). Training data were sampled 30 times, each time extracting 15% of the comprehensive training dataset. Each selected sample was used to create a tree model resulting in an ensemble of 30 models. These models were applied to classify each of the 23 annual AVHRR data composites. The 30 classification results were then combined by averaging to estimate the probability of burning from 0 to 100%. Based on the probability, a simple majority threshold was used to tag the pixel as burned. Lastly, we used a 3 x 3 majority kernel to filter and smooth the classification results. This filtering may have removed some correctly mapped small fires, but we employed it to ensure conservative estimation of burned area and lower commission errors.

## **Validation**

### *Burned area map validation*

For the validation of the results we created an additional dataset of burned areas from higher resolution imagery (TM, ETM+, MODIS, MSU-SK). Red-green-blue composites were created for each image, using bands 2 (841-876 nm), and 1 (620-670 nm) from MODIS, bands 5 (1550-1750 nm), 4 (760-900 nm), and 3 (630-690nm) from TM/ETM+, and bands 4 (810-1000

nm), 2 (600-720 nm), and 1 (540-600 nm) from MSU-SK imagery. We validated our classifications for each year for which validation data were available.

Similarly to training data selection, validation data were obtained by visual image interpretation, shown to be at least as precise as automatic methods (Bowman et al. 2002). Delineating burned areas was straightforward due to distinct fire scar boundaries, often confined by linear features such as roads (Figure 7). The easy recognition of fires was further facilitated by the lack of other disturbances in the area that could have resulted in comparable patterns. In many cases fire scars remained visible for 2 to 3 years after burning, unless a new fire overrode an older scar (Figure 7). The elevated brightness of burned areas was also quite stable intra-annually (Figure 4). The validation dataset was rasterized to the 1.1 x 1.1 km pixels. Omission and commission errors were calculated for each year by combining binary burned classification and rasterized validation datasets. Producer's and user's accuracies were calculated from the four possible combinational classes (Congalton and Green 1998). Additionally, we calculated the Kappa coefficient, which is a proportion of agreement obtained after removing the proportion of agreement expected to occur by chance (Congalton and Green 1998).

#### *Out of season validation*

Because the AVHRR image dates were not uniform, we also checked if sub-optimal image dates caused omission of some fires. We tested this by comparison with numbers of active fire hotspots captured by both TERRA and AQUA satellites (MOD14A1 and MYD14A1). Based on the active fire data, we calculated the percentage of fires between August 20<sup>th</sup> and August 30<sup>th</sup> compared to the entire month of August, and the percentage of fires in August and September to the entire fire season for each year from 2000 to 2007. Though the number of active fire detections is correlated with the size of the burned area, this measure is not always reliable. Since

some of the selected AVHRR images were collected in early August and we found some active fires in September that were not covered by the selected images, we extended the validation dataset until the end of September for the years 2002 to 2007. This allowed us to estimate how much burned area was missed due to the lack of optimally timed AVHRR images in some years.

#### *Model transferability validation*

Due to the coarse resolution of AVHRR satellite imagery, the visual estimation of training data may not always be reliable. Furthermore, in some years, no burned areas could be found for training. Using a comprehensive training dataset and a uniform set of decision tree models for all years captured the variability in burned area signals, and ensured that burning was not simply missed in a given year when visually interpreting the AVHRR data. However, the use of the comprehensive training dataset needed validation. To explore the transferability of the model to years for which there was no burned area training data, we removed training data for a set of years where amount of burning exceeded 15,000 km<sup>2</sup> (2000, 2002, and 2005 to 2007) from the comprehensive training dataset and repeated the classification. These tests quantified the extent to which uniform set of models was able to detect burning in one year in the absence of samples from this year in the comprehensive training set.

## **Results**

### **Annual burned area**

Burned area increased dramatically from virtually no burning in the 1980s and the first half of the 1990s, to up to 19% of the study area (3,600 km<sup>2</sup>) burned annually after 2000 (Figure 5 and Table 1). Among the 23 years that we studied, nine had substantial amounts of burned

areas, while the other 14 had less than 2% of the area burned. Eight out of these nine high-burn years occurred after 1998 (Table 1). Large-scale burning started abruptly in 1997 and 1998, when 5% of the territory was burned and has continued regularly until today. However, burned area showed a high degree of variability, with one or two years of burning often followed by a year with negligible burned area.

Among the different AVHRR-derived bands that we used for the classification, the most important were the NDVI of the fall image and the difference between spring and fall values of band 1 and band 3. These metrics explained 47% of the variability; each contributed respectively 24%, 14%, and 9% to the decrease of deviance for all 30 decision tree models. Other metrics were equally valuable and contributed on average 6% of the deviance decrease.

Large fires contributed to the majority of the burned area. As estimated using MODIS data, burns larger than 250 km<sup>2</sup> represented 48% of total burned area. Though small fires (less than 60 km<sup>2</sup>) represented 77% of the total number of fires since 2000, their share accounted only for 18% of total burned area (Figure 8). On average, the five largest fires in a given year represented 66% of the total burned area for that year. Since 2000, there was also a marked increase in the area of the largest fire, which grew from 58% of the total area burned in 2000 to almost 70% in 2006, and 71% in 2007 (1,930, 3,740, and 3,120 km<sup>2</sup> respectively).

## Validation

### *Burned area map validation*

Validation of the burned area maps relied on data from different satellite sensors. For burned area delineations we used a total of 1,066 MODIS images (a mean of 133 images per year, range 71-195), and 10 RESURS images (mean 3, range 2-4). MODIS data was available for the entire burning season; RESURS data also covered the entire burning season, except 1998, where

the last RESURS image was acquired on August, 20. Validation with Landsat data was based on September or late August imagery, and we used only one late season image per year. Average producer's accuracy of the burned area class in years where at least 200 km<sup>2</sup> burned was 68% (Table 1). Average user's accuracy for the same years was 56% (Table 1). The Kappa coefficient for the same year was 52% or moderate, according to classification scale by Congalton (1996). It was not possible to validate independently 10 out of 23 years (1985, 1990-1995) due to the lack of MODIS, Landsat, and RESURS data. No signs of burned areas were found for 7 of the validated years (1986, 1988, 1996, 1999, and 2003). Official statistics, available since 2002, were well correlated with the estimated trend, but showed generally much less burned area than our classifications (Figure 5).

#### *Temporal distribution and out of season validation*

We estimated the temporal distribution of fires for 2000 to 2007 using MODIS active fire hotspots and our burned areas boundaries derived for validation. As reported previously (Carmona-Moreno et al. 2005) burning predominantly started in June and ended in August (Figure 9). Few active fire hotspots were detected before June and in late August and September. The percentage of active fire counts in August compared to a 3-month total was 26% (32% annual average), and only 9% (12% annual average) of the total number of active fires were detected in September. In August, most of the burning happened in the beginning and the middle of the month. The percentage of active fires detected during the last ten days of August was only 12% of the total number of active fire counts during August (14% annual average).

Burn areas mapped using MODIS data that occurred after the AVHRR images were taken showed that only a small proportion of burns were missed due to the timing of images. With the exception of 2003, less than 2% of total burned area occurred after the AVHRR images



were taken (20 to 90 km<sup>2</sup> in a given years). However, in 2003 only 3 fires occurred in total, and one of those 3 occurred after the AVHRR image date and covered 70 km<sup>2</sup> or 63% of the total burned area in that year. Some burning may have taken place outside the windows captured by the AVHRR in those years for which no MODIS data were available. However, field observations, expert opinion and examination of imagery for period before and after fire season indicated that such burning was overall very limited.

### *Model transferability*

We also tested how well our pooled training data could classify burns in years for which no training data were available (or withheld to test the transferability of the training data). Overall, the transferability was moderate. The amount of burned area for a particular year detected without training data from this year ranged from 84% (2002), 52% (2006), 44% (2007), down to as low as 17% (2005), of the area predicted with a full sample. Transferred models were successful in detecting burns that led to the complete removal of vegetation resulted in big areas of bare ground. Poor transferability in 2005 was likely due to a somewhat abnormal fire season. Unlike other test years, most of the burned areas in 2005 happened late in the year and showed low reflectance in visible bands due to large amounts of remaining soot.

## ***Discussion***

Our goals were twofold. First, we aimed to estimate changes in burned area before and after the precipitous declines in livestock caused by socioeconomic changes following the breakdown of the Soviet Union. Second, we wanted to develop a method to map burned areas from AVHRR data suitable for the parts of the world for which only data from CLASS were available. Both goals were met successfully.

The main finding from our study was a dramatic increase in burned areas in the arid grasslands of Southern Russia starting in the late 1990s (Figure 5). The area burned each year jumped from almost no fires in the 1980s to large-scale burning, covering up to 20% of study area in a single year after the mid-1990s. Since 1998, on average 1,381 km<sup>2</sup> burned per year (9% of the study area, Figure 10). At this rate, the entire area will burn every 11 years. Fires occurred almost exclusively during the driest season of the year (Carmona-Moreno et al. 2005). Although no data on lightning occurrences and dry thunderstorms existed for the area, our field experience suggested that they are rare. Thus we assume that most of the fires are human-caused, resulting from transportation, mainly to and from local herding enterprises, hunting activities, including illegal poaching for indigenous saiga antelope (*Saiga tatarica*), and carelessness (e.g., wide use of old machinery, smoking, etc.). There is no history of grassland burning for pasture management in the study area (Y. Arylov, personal communication).

### **Burned area trends, vegetation, grazing, and socioeconomic changes**

The abrupt increase in burned areas that we observed followed a sharp decrease in livestock abundance around the time of breakdown of Soviet Union in 1991. We suggest that the main mechanism leading to the increase in fires was the removal of livestock, predominantly sheep. Reduced disturbance by livestock allowed vegetation to recover, and caused a gradual increase in biomass and fuel connectivity. Grass-dominated vegetation communities (*Stipa* spp.) have increased particularly rapidly (Neronov 1998). Our results suggest that this recovery of grasslands led to an increase of the amount of dry litter that permitted the consequent significant increase in burning.

Increase in burned areas was also concomitant with the reduction of fire suppression activities in Russia, resulting from economic hardship after the breakdown of Soviet Union in

1991, and the major economic crisis of 1998 (Shvidenko and Goldammer 2001). Unemployment caused an increase in the number of visits by locals to the steppe in search of other sources of income. Burning was used to assist poaching of the endangered saiga antelope and led to near obliteration of the last population of saiga in European Russia (Lushchekina and Struchkov 2001). Official burned areas statistics were only available after 2002. The trends in the burned area statistics correlated well with our AVHRR-based estimates, but official statistics highly underestimated the area burned. Underestimation of fires in governmental data is due to the fact that only fires that were actively suppressed were included in official statistics (Soja et al. 2004).

The strong increase of annual burned area in late 1990s in our study region is similar to other parts of the grasslands of Central Asia and Mongolia. Being part of the Soviet block, these countries showed similar socioeconomic changes and declines in livestock numbers. Though fire data are scarce for Central Asia, there has been a 6-fold increase in steppe and forest fires in Mongolia between the periods of 1985-1995 and 1996-1997 (Erdenesaikhan and Erdenetuya 1999; Velsen-Zerweck 2002), similar to the pattern found in our study.

The area that burned most often occurred in a strictly-protected nature reserve (Chernye Zemli Zapovednik) established in 1990, which obtained Biosphere Reserve status in 1993, and covers 1,220 km<sup>2</sup> (6% of the study area). The reserve allows research, conservation efforts, and education, but prohibits any other human activities, like livestock grazing or haymaking. The extra protection may have facilitated fuel buildup in the protected area. Together with the livestock declines in the surrounding areas, this may have caused increasing numbers of fires of medium and large size. Although there is no 'let burn' policy in Russia's protected areas, according to our estimation, the reserve burned almost entirely in 2000, 2002, 2006, and 2007, most likely due to the lack of funds for effective fire suppression.

## Satellite data limitations

The only satellite dataset with annual data records spanned from 1985 until present that was available for southern Russia was CLASS-type AVHRR data. Despite the limitations of CLASS-type AVHRR data, we were able to develop a method to map burned areas and to detect long-term trends in annual burned area. However, our accuracy assessment highlighted the limitations of the available coarse-resolution satellite data. We could not use burned area mapping algorithms that require multiple ( $>2$ ) images for a given year due to the lack of good data in the archives, excessive cloud cover, as well as geometric and radiometric artifacts. In our study area, daily AVHRR data that could be composited were not collected by local stations until the late 1990s. We suggest that the lack of data is largely responsible for the reported classification accuracies, and higher accuracies could have been obtained with if daily acquisitions were available. However, despite its limitations, AVHRR data often provides the only means possible to reconstruct dense time series of burned area estimates for long time spans, prior to the launch of MODIS in 2000. Other remote sensing data, while providing better spectral and spatial resolution, lack temporal resolution, spatial coverage, and the long-term data record necessary to give clear answer to the question of how burned area changed throughout the recent 20+ years. This is especially true for those regions of the world where remote sensing data records are sparse due to poor archiving or lack of acquisitions. Recent developments in remote sensing data distribution policy, such as the ‘freeing’ of the Landsat data archives, may make it possible to map burned areas with high resolution in time and space. However, we caution that many areas of the world have acquisition gaps, and methods like the one we present here will be needed to estimate burned area trends for long time series.

## **Validation of the burned area estimates and the burned area mapping approach**

Our study dealt with a particularly long time series of satellite observations and various satellite sources were used for validation based on their availability. Though the accuracy of the estimates derived in this study was moderate, similar accuracies were found in comparable environments by studies using better data and more advanced methodology (Roy and Boschetti 2009). Differences among the imagery used for validation may have resulted in some inconsistencies in the reported accuracy rates, which we could not quantify. Generally, the dataset with highest temporal frequency (i.e., MODIS) was best suited to delineation burned areas for our validation dataset, followed by less frequent, but spatially more detailed Landsat/TM, and finally the only data available for late 1990s (i.e., RESURS/MSU-SK)..

Our analysis of the fire season with MODIS active fire data showed that a few fires could have occurred after the dates of our AVHRR images, but burned areas potentially missed were in all likelihood negligible. Our conclusion that fires were essentially absent before 1998 was also supported by our test of the transferability of the decision trees to years for which no training data for burned areas were available. On average of 42% of the burned areas detected with full training data were still detected when training data for a given year were omitted. That suggests that even if our visual interpretation of fires in the AVHRR data prior to 1998 missed fires erroneously, our models would have still captured a considerable portion of them.

We found that the largest fires captured up to 70% of the total area burned in a given year (Figure 6). Burned area classifications in Southern California (Minnich 1983), the Intermountain West (Knapp 1998), Mongolia (Erdenesaikhan and Erdenetuya 1999) and Australia (Yates et al. 2008) found similar pattern with the most of the burned area attributed to few large contiguous

burns. We caution though, that the share of large burned areas may have been overestimated due to the low resolution bias of AVHRR data, resulting from the 1.1 x 1.1 km pixel size (Boschetti et al. 2004), and our use of a majority filter. Omission of small burned areas with AVHRR data is especially critical in mosaic environments representative of crop production systems (Laris 2005). We cannot rule out that fires were just as frequent prior to 1998, but of such small size that AVHRR missed them all. However, this explanation for the observed increase in burned area is highly unlikely. First, size class analysis with higher-resolution MODIS data during the years since 2000, when fires were widespread, showed a clear dominance of large fires (Figure 8), supported by the contiguous nature of grassland vegetation and the absence of crop production. And second, comparable rates of burning expressed as much smaller fires would require much more dense population (to provide ignitions, but also to suppress fires before they got large) or fine patterns of land use (to prevent fires from getting large). Neither was present in the study area historically.

Lastly, our finding that there was essentially no fire prior to 1998 is supported by the only other remote sensing study in the study area. A change detection for 1989 to 1998 by Hoelzel (2002) did not mention any burned areas as well. Though not quantified, sudden and vast increase in burning in our study area after the mid-1990s has also been mentioned in other studies (Shilova et al. 2007) and is corroborated by members of local communities and other scientists working in the area (Bakinova T.I., Dzhapova R.R., Neronov V.V., personal communications).

## **Concluding remarks**

Our study provides one of the first estimates of burned area trends for the arid grasslands of Central Asia, and represents one of the longest burn area time series obtained using remote

sensing here and elsewhere in the world. Drastic socioeconomic changes often lead to changes of a particular disturbance agent. In arid grasslands, the decline in grazing might result in the transition of the ecosystem to a different state. However, changes in vegetation due to reduced grazing likely lead to the substitution of grazing by fire as the main disturbance. Interestingly, the replacement of the disturbance agents did not occur immediately, but exhibited a time lag. Knowing the time lag between disturbances is important in order to make accurate predictions, and we speculate that fire may again decline given continuing restoration of livestock numbers returning to their historic highs (Fig. 1). Whether we are seeing recovery of the ecosystem in the long term or transition to a new state remains a question.

## ***Acknowledgements***

We gratefully acknowledge support for this research by the NASA Land-Cover and Land-Use Change Program, the National Geographic Society, and by a NASA Earth System Science Dissertation Fellowship for M. Dubinin (07-Earth07F-0053). We also thank the R&D Center Scanex (Moscow, Russia) for providing RESURS imagery.

## ***Bibliography***

- Akiyama, T., & Kawamura, K. (2007). Grassland degradation in China: Methods of monitoring, management and restoration. *Grassland Science*, 53, 1-17
- Archer, S., Schimel, D.S., & Holland, E.A. (1995). Mechanisms of shrubland expansion - land-use, climate or CO<sub>2</sub>. *Climatic Change*, 29, 91-99
- Archibald, S., & Bond, W.J. (2004). Grazer movements: spatial and temporal responses to burning in a tall-grass African savanna. *International Journal of Wildland Fire*, 13, 377-385
- Arino, O., Piccolini, I., Kasischke, E., Siegert, F., Chuvieco, E., Martin, M.P., Li, Z., Fraser, R., Eva, H., Stroppiana, D., Pereira, J., Silva, J.M.N., Roy, D., & Barbosa, P.M. (2001). Methods of mapping burned surfaces in vegetation fires. In F.J. Ahern, J.G. Goldammer & C.O. Justice (Eds.) *Global and regional vegetation fire monitoring from space: planning a coordinated international effort* (pp. 227-255). The Hague SPB Academic Publishing
- Bahre, C.J. (1991). *A Legacy of Change: historic human impact on vegetation of the Arizona borderlands*. Tucson University of Arizona Press
- Barbosa, P.M., Gregoire, J.-M., & Pereira, J.M.C. (1999). An Algorithm for Extracting Burned Areas from Time Series of AVHRR GAC Data Applied at a Continental Scale. *Remote Sensing of Environment*, 69, 253-263
- Bond, W.J., & van Wilgen, B.W. (1996). *Fire and plants*. London: Chapman and Hall
- Boschetti, L., Flasse, S.P., & Brivio, P.A. (2004). Analysis of the conflict between omission and commission in low spatial resolution dichotomic thematic products: The Pareto Boundary. *Remote Sensing of Environment*, 91, 280-292
- Bowman, D., Zhang, Y., Walsh, A., & Williams, R.J. (2002). Experimental comparison of four remote sensing techniques to map tropical savanna fire-scars using Landsat-TM imagery. In,



- Conference on Fire and Savanna Landscapes in Northern Australia* (pp. 341-348). Darwin, Australia
- Breiman, L. (1984). *Classification and regression trees*. Belmont, Calif.: Wadsworth International Group
- Breiman, L. (1996). Bagging predictors. *Machine Learning*, 24, 123-140
- Brooks, K., & Gardner, B. (2004). Russian agriculture in the transition to a market economy. *Economic Development and Cultural Change*, 52, 571-586
- Brooks, M.L., D'Antonio, C.M., Richardson, D.M., Grace, J.B., Keeley, J.E., DiTomaso, J.M., Hobbs, R.J., Pellant, M., & Pyke, D. (2004). Effects of invasive alien plants on fire regimes. *Bioscience*, 54, 677-688
- Burcher, C.L., Valett, H.M., & Benfield, E.F. (2007). The land-cover cascade: Relationships coupling land and water. *Ecology*, 88, 228-242
- Carmona-Moreno, C., Belward, A., Malingreau, J.P., Hartley, A., Garcia-Alegre, M., Antonovskiy, M., Buchshtaber, V., & Pivovarov, V. (2005). Characterizing interannual variations in global fire calendar using data from Earth observing satellites. *Global Change Biology*, 11, 1537-1555
- Chuvieco, E., Englefield, P., Trishchenko, A.P., & Luo, Y. (2008). Generation of long time series of burn area maps of the boreal forest from NOAA-AVHRR composite data. *Remote Sensing of Environment*, 112, 2381-2396
- CIESIN, & CIAT (2005) Gridded Population of the World Version 3 (GPWv3): Population Density Grids. Palisades, NY; Available at <http://sedac.ciesin.columbia.edu/gpw>. Last accessed: 06.28.2009: Socioeconomic Data and Applications Center (SEDAC), Columbia University

- CIESIN, FAO, & CIAT (2005) Gridded Population of the World: Future Estimates, 2015 (GPW2015): Population Density Grids. Palisades, NY; Available at <http://sedac.ciesin.columbia.edu/gpw>. Last accessed: 06.28.2009: Socioeconomic Data and Applications Center (SEDAC), Columbia University
- Congalton, R.G., & Green, K. (1998). *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices*: CRC Press
- Csiszar, I., Denis, L., Giglio, L., Justice, C.O., & Hewson, J. (2005). Global fire activity from two years of MODIS data. *International Journal of Wildland Fire*, 14, 117-130
- Csiszar, I., Justice, C.O., Mcguire, A.D., Cochrane, M.A., Roy, D.P., Brown, F., & al., e. (2004). Land use and fires In G.A. Gutman & et al (Eds.) *Land change science: Observing, monitoring, and understanding trajectories of change on the earth's surface*. Kluwer Academic Publishers
- Diamond, J. (2001). Ecology: Dammed Experiments! *Science*, 294, 1847-1848
- Erdenesaikhan, N., & Erdenetuya, M.M. (1999). Forest and steppe fire monitoring in Mongolia using satellite remote sensing. *International Forest Fire News*, 21, 71-74
- Friedl, M.A., & Brodley, C.E. (1997). Decision tree classification of land cover from remotely sensed data. *Remote Sensing of Environment*, 61, 399-409
- Giglio, L., van der Werf, G.R., Randerson, J.T., Collatz, G.J., & Kasibhatla, P. (2006). Global estimation of burned area using MODIS active fire observations. *Atmospheric Chemistry and Physics*, 6, 957-974
- Golub, V.B. (1994). The desert vegetation communities of the Lower Volga Valley. *Feddes Repertorium*, 105, 499-515

Gong, P., Pu, R.L., Li, Z.Q., Scarborough, J., Clinton, N., & Levien, L.M. (2006). An integrated approach to wildland fire mapping of California, USA using NOAA/AVHRR data.

*Photogrammetric Engineering and Remote Sensing*, 72, 139-150

Hansen, M.C., Defries, R.S., Townshend, J.R.G., & Sohlberg, R. (2000). Global land cover classification at 1km spatial resolution using a classification tree approach. *International Journal of Remote Sensing*, 21, 1331-1364

Hawbaker, T.J., Radeloff, V.C., Syphard, A.D., Zhu, Z.L., & Stewart, S.I. (2008). Detection rates of the MODIS active fire product in the United States. *Remote Sensing of Environment*, 112, 2656-2664

Hoelzel, N., Haub, C., Ingelfinger, M.P., Otte, A., & Pilipenko, V.N. (2002). The return of the steppe: Large-scale restoration of degraded land in southern Russia during the post-Soviet era. *Journal for Nature Conservation (Jena)*, 10, 75-85

Justice, C.O., Vermote, E., Townshend, J.R.G., Defries, R., Roy, D.P., Hall, D.K., Salomonson, V.V., Privette, J.L., Riggs, G., Strahler, A., Lucht, W., Myneni, R.B., Knyazikhin, Y., Running, S.W., Nemani, R.R., Wan, Z.M., Huete, A.R., van Leeuwen, W., Wolfe, R.E., Giglio, L., Muller, J.P., Lewis, P., & Barnsley, M.J. (1998). The Moderate Resolution Imaging Spectroradiometer (MODIS): Land remote sensing for global change research. *IEEE Transactions on Geoscience and Remote Sensing*, 36, 1228-1249

Knapp, P.A. (1998). Spatio-temporal patterns of large grassland fires in the Intermountain West, USA. *Global Ecology and Biogeography*, 7, 259-272

Kroonenberg, S.B., Rusakov, G.V., & Svitoch, A.A. (1997). The wandering of the Volga delta: A response to rapid Caspian sea-level change. *Sedimentary Geology*, 107, 189-209

- Kučera, J., Yasuoka, Y., & Dye, D.G. (2005). Creating a forest fire database for the Far East of Asia using NOAA/AVHRR observation. *International Journal of Remote Sensing*, 26, 2423-2439
- Kuemmerle, T., Hostert, P., Radeloff, V.C., Perzanowski, K., & Kruhlov, I. (2007). Post-socialist forest disturbance in the Carpathian border region of Poland, Slovakia, and Ukraine. *Ecological Applications*, 17, 1279-1295
- Kurinova, Z.S., & Belousova, Z.N. (1989). *Effective use of forage resources*. Elista: Kalmyk book publishing
- Lambin, E.F., & Geist, H. (2006). *Land-use and land-cover change: local processes and global impacts*: Springer
- Laris, P.S. (2005). Spatiotemporal problems with detecting and mapping mosaic fire regimes with coarse-resolution satellite data in savanna environments. *Remote Sensing of Environment*, 99, 412-424
- Lee, D.S., Storey, J.C., Choate, M.J., & Hayes, R.W. (2004). Four years of Landsat-7 on-orbit geometric calibration and performance. *IEEE Transactions on Geoscience and Remote Sensing*, 42, 2786-2795
- Liedloff, A.C., Coughenour, M.B., Ludwig, J.A., & Dyer, R. (2001). Modelling the trade-off between fire and grazing in a tropical savanna landscape, northern Australia. *Environment International*, 27, 173-180
- Lushchekina, A.A., & Struchkov, A. (2001). The saiga antelope in Europe: once again on the brink? *The Open Country*, 3, 11-24
- Mack, R.N., & Thompson, J.N. (1982). Evolution in steppe with few large, hooved mammals. *The American Naturalist*, 119, 757-773

- Maggi, M., & Stroppiana, D. (2002). Advantages and drawbacks of NOAA-AVHRR and SPOT-VGT for burnt area mapping in a tropical savanna ecosystem. *Canadian Journal of Remote Sensing*, 28, 231-245
- Meyn, A., White, P.S., Buhk, C., & Jentsch, A. (2007). Environmental drivers of large, infrequent wildfires: the emerging conceptual model. *Progress in Physical Geography*, 31, 287-312
- Miettinen, J., Langner, A., & Siegert, F. (2007). Burnt area estimation for the year 2005 in Borneo using multi-resolution satellite imagery. *International Journal of Wildland Fire*, 16, 45-53
- Minnich, R.A. (1983). Fire Mosaics in Southern California and Northern Baja California. *Science*, 219, 1287-1294
- Neronov, V.V. (1998). Antropogenic grasslands encroachment of desert pastures of North-Western Precaspian plains. *Successes of Modern Biology*, 118, 597-612 (in Russian)
- Pereira, J.M.C. (1999). A comparative evaluation of NOAA/AVHRR vegetation indexes for burned surface detection and mapping. *IEEE Transactions on Geoscience and Remote Sensing*, 37, 217-226
- Pu, R.L., Li, Z.Q., Gong, P., Csiszar, I., Fraser, R., Hao, W.M., Kodragunta, S., & Weng, F.Z. (2007). Development and analysis of a 12-year daily 1-km forest fire dataset across North America from NOAA/AVHRR data. *Remote Sensing of Environment*, 108, 198-208
- Pyne, S.J. (1984). *Introduction to Wildland Fire*. New York: John Wiley and Sons
- Razafimpanilo, H., Frouin, R., Iacobellis, S.F., & Somerville, R.C.J. (1995). Methodology for estimating burned area from AVHRR reflectance data. *Remote Sensing of Environment*, 54, 273-289

ROSSTAT (2003). Agriculture in Republic of Kalmykia. *Kalmykia in numbers (in Russian)*.

Elista

ROSSTAT (2007). Agriculture in Republic of Kalmykia. *Kalmykia in numbers (in Russian)*.

Elista

Roy, D.P., & Boschetti, L. (2009). Southern Africa Validation of the MODIS, L3JRC, and GlobCarbon Burned-Area Products. *IEEE Transactions on Geoscience and Remote Sensing*, 47, 1032-1044

Saiko, T., & Zonn, I. (1997). Europe's first desert. In M. Glantz & I. Zonn (Eds.) *Scientific, Environmental and Political Issues in the Circum-Caspian Region* (pp. 141-144). Dordrecht, Boston and London Kluwer Academic Publishers

Shilova, S.A., Neronov, V.V., Kasatkin, M.V., Savinetskaya, L.E., & Chabosvsky, A.V. (2007). Modern burning in semi-deserts of Southern Russia: influence on vegetation and rodents populations. *Uspekhi Sovremennoi Biologii*, 127, 372-386

Shvidenko, A., & Goldammer, J.G. (2001). Fire situation in Russia. *International Forest Fire News*, 24, 41-59

Soja, A.J., Sukhinin, A.I., Cahoon, D.R., Shugart, H.H., & Stackhouse, P.W. (2004). AVHRR-derived fire frequency, distribution and area burned in Siberia. *International Journal of Remote Sensing*, 25, 1939-1960

Song, C., Woodcock, C.E., Seto, K.C., Lenney, M.P., & Macomber, S.A. (2001). Classification and change detection using Landsat TM data: When and how to correct atmospheric effects? *Remote Sensing of Environment*, 75, 230-244

Stohl, A., Berg, T., Burkhardt, J.F., Fjaeraa, A.M., Forster, C., Herber, A., Hov, O., Lunder, C., McMillan, W.W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Strom, J.,

- Torseth, K., Treffeisen, R., Virkkunen, K., & Yttri, K.E. (2007). Arctic smoke - record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006. *Atmospheric Chemistry and Physics*, 7, 511-534
- Stroppiana, D., Tansey, K., Gregoire, J.M., & Pereira, J.M.C. (2003). An algorithm for mapping burnt areas in Australia using SPOT-VEGETATION data. *IEEE Transactions on Geoscience and Remote Sensing*, 41, 907-909
- Sukhinin, A.I., French, N.H.F., Kasischke, E.S., Hewson, J.H., Soja, A.J., Csiszar, I.A., Hyer, E.J., Loboda, T., Conrad, S.G., Romasko, V.I., Pavlichenko, E.A., Miskiv, S.I., & Slinkina, O.A. (2004). AVHRR-based mapping of fires in Russia: New products for fire management and carbon cycle studies. *Remote Sensing of Environment*, 93, 546-564
- Tansey, K., Gregoire, J.M., Stroppiana, D., Sousa, A., Silva, J., Pereira, J.M.C., Boschetti, L., Maggi, M., Brivio, P.A., Fraser, R., Flasse, S., Ershov, D., Binaghi, E., Graetz, D., & Peduzzi, P. (2004). Vegetation burning in the year 2000: Global burned area estimates from SPOT VEGETATION data. *Journal of Geophysical Research-Atmospheres*, 109
- van der Werf, G.R., Randerson, J.T., Giglio, L., Collatz, G.J., Kasibhatla, P.S., & Arellano, A.F. (2006). Interannual variability in global biomass burning emissions from 1997 to 2004. *Atmospheric Chemistry and Physics*, 6, 3423-3441
- Velsen-Zerweck, M.v. (2002). Socio-economic causes of forest loss in Mongolia. *Sozialökonomische Schriften zur Ruralen Entwicklung* (p. xxi + 357 pp.)
- Vinogradov, B.V. (1995). Forecasting dynamics of deflated sands of Black Lands, Kalmykia using aerial photography. In N. Zonn I.S., V.M. (Ed.) *Biota and Environment of Kalmykia* (pp. 259-268). Moscow, Elista (in Russian)

- Walter, H., & Box, E. (1983). Overview of Eurasian continental deserts and semideserts. *Ecosystems of the World* (pp. 3-269). Amsterdam Elsevier
- Yates, C.P., Edwards, A.C., & Russell-Smith, J. (2008). Big fires and their ecological impacts in Australian savannas: size and frequency matters. *International Journal of Wildland Fire*, 17, 768-781
- Zhang, K., Yu, Z., Li, X., Zhou, W., & Zhang, D. (2007). Land use change and land degradation in China from 1991 to 2001. *Land Degradation & Development*, 18, 209-219
- Zonn, S.V. (1995). Desertification of nature resources in Kalmykia during last 70 years and protection measures. *Biota and Environment of Kalmykia* (pp. 12-52). Moscow, Elista (in Russian)



## Tables

Table 1. Annual burned area estimation and validation results for study period.

<b>Year</b>	<b>Burned area, km<sup>2</sup></b>	<b>Burned area % total</b>	<b>Producer's accuracy, burned</b>	<b>User's accuracy, burned</b>	<b>Producer's accuracy, unburned</b>	<b>User's accuracy, unburned</b>	<b>Kappa</b>
1985	0	0.0	N/A	N/A	100.0%	100.0%	N/A
1986	25	0.1	0.0%	N/A	100.0%	99.9%	0.0%
1987	0	0.0	0.0%	N/A	100.0%	98.7%	N/A
1988	74	0.4	0.0%	N/A	100.0%	99.6%	0.0%
1989	21	0.1	0.0%	N/A	100.0%	99.9%	0.0%
1990	4	0.0	N/A	N/A	100.0%	100.0%	0.0%
1991	0	0.0	N/A	N/A	100.0%	99.5%	N/A
1992	266	1.4	N/A	N/A	100.0%	98.6%	0.0%
1993	90	0.5	N/A	N/A	100.0%	99.5%	0.0%
1994	0	0.0	N/A	N/A	100.0%	100.0%	N/A
1995	0	0.0	N/A	N/A	100.0%	99.8%	N/A
1996	0	0.0	0.0%	N/A	100.0%	99.4%	N/A
1997	518	2.7	18.5%	82.3%	99.9%	97.8%	29.4%
1998	933	4.9	92.5%	34.2%	90.8%	99.6%	46.1%
1999	0	0.0	0.0%	N/A	100.0%	99.9%	N/A
2000	1,602	8.5	78.2%	48.9%	92.4%	97.9%	55.5%
2001	417	2.2	62.3%	28.4%	96.4%	99.1%	37.1%
2002	2,586	13.7	82.6%	64.1%	92.7%	97.1%	67.2%
2003	34	0.2	28.2%	0.4%	82.5%	99.8%	1.7%
2004	567	3.0	67.8%	32.5%	95.6%	99.0%	41.4%
2005	2,193	11.6	52.0%	63.5%	96.1%	93.8%	50.5%

2006	3,781	20.0	86.5%	73.8%	92.3%	96.5%	74.2%
2007	4,397	23.3	72.8%	75.4%	92.8%	91.8%	66.4%

Table 2. Detection rates of burned areas in different size class categories.

	Burned area patch size (km <sup>2</sup> )							
	<10	10-20	20-35	35-60	60-100	100-150	150-250	>250
Number of fires	144	64	48	39	31	19	16	14
Number of fires detected	31	27	23	30	24	16	16	13
Percent detected	22%	42%	48%	77%	77%	84%	100%	93%

## Figures

Figure 1 Total sheep population in three administrative regions of the Republic of Kalmykia (Chernozemelsky, Iki-Burulsky, Lagansky) which closely correspond to the boundaries of our study area (ROSSTAT 2003, 2007).




Figure 2. Study area location (hatched polygon with black outline, grey line – Republic of Kalmykia; the grayscale background represents shaded relief (SRTM30)).



Figure 3. Feathergrass (*Stipa* sp.) (top) and sage (*Artemisia* sp.) (bottom) dominated communities in the beginning (left) and the end (right) of the vegetation season.



Figure 4. 16-day MODIS reflectance values in bands 1 (triangles) and 2 (squares) for a sample area (median of 9 pixels) affected by fire (hatched line denotes the fire date). Winter-time snow contaminated values are not shown.




Figure 5. Estimated burned area according to the AVHRR analysis (solid line), validation dataset (dotted line, triangles) and official burned area data (dashed line, squares).




Figure 6. An example of AVHRR-based burned area map for year 2000 (red polygons) comparing with MODIS-derived burned area validation data (outlined in black) shown on top of the AVHRR image from Aug 1, 2000. The boundary of analysis region is shown in magenta.






Figure 7. Example of before-burning and after-burning in a Landsat TM image of September, 1988 and an ETM+ image of July, 2001 (5-4-3 band combination). Ubiquitous fire-scars bounded by roads in the 2001 image, most of these scars are from previous years (marked with arrows). The photo represents an area burned in September, 4, 2005, photo taken March, 18, 2006




Figure 8. Number (line) and area (bars) of burn areas from 2000 to 2007 summarized by size class.




Figure 9. Fire seasonality, i.e., the burned area by month, based on the MODIS image interpretation.





Figure 10. Burned area mapped for selected years with high fire activity. Gray shading – AVHRR-based burned area maps, black outlines – validation burned area maps (1998 – RESURS-based, others – MODIS-based).



## **Chapter 2: Relationships between climate, livestock, and fire in arid ecosystems of Southern Russia**

### ***Abstract***

Fire is an important natural disturbance process in arid grasslands but current fire regimes are largely the result of both human and natural processes and their interactions. The breakdown of the Soviet Union in 1991 spurred substantial socioeconomic changes and was ultimately followed by a rapid increase in burned area in southern Russia. It is whether increase in burned area was caused by decreasing livestock numbers, vegetation changes, climate change, or interactions of these factors. Our research goal was to identify the driving forces behind the increase in burned area in the arid grasslands of southern Russia. Our study area encompassed 19,000 km<sup>2</sup> in the Republic of Kalmykia in southern Russia. We analyzed annual burned area from 1985 to 2006 as function of livestock population, NDVI, precipitation, temperature, and broad-scale oscillation indices using best subset regression and structural equation modeling. Our results supported the hypothesis that the vegetation recovered within 5-6 years after the livestock declined in the beginning of the 1990s, to a point at which large fires could be sustained. Climate was an important explanatory factor for burning, but mainly after 1996 when lower livestock numbers allowed fuels to accumulate. Ultimately, our results highlight the complexity of coupled human-natural systems, and provide an example of how abrupt socioeconomic change may affect fire regimes.

## ***Introduction***

Fire is one of the main disturbance agents in grasslands and savannahs, and can be an indicator of ecosystem change. Fire shapes vegetation structure and composition and represents an important land-management tool (Pyne, 1984). Grasslands, woody savannahs, and savannahs contain more than 60% of the annual global burned area (Tansey et al., 2004), and in regions with high aridity, such as Central Asia, grassland fires account for 80% of all active fire counts (Csiszar et al., 2005). Fuel loads and emissions from grassland burning are relatively small (van der Werf et al., 2006), but grassland fires can foster the spread of invasive species (Brooks et al., 2004), affect wildlife habitat (Archibald and Bond, 2004), and generate air pollution that can spread as far as the Arctic (Stohl et al., 2007). Furthermore, interactions between grassland fires and human land use may lead to ecosystem degradation, hydrologic changes, soil disturbance, and shrub encroachment (Archer et al., 1995). At the same time, fire is an indispensable component of grassland ecosystems because fires stimulate germination and regrowth (Letnic, 2004), and hence fires are often set deliberately to enhance pastures and raise the availability of nutritious forage for livestock (Guevara et al., 1999). However, management, conservation, and restoration of arid grasslands require an understanding of the driving forces of burning, and such an understanding can only be gained from long-term fire records. Our goal was to understand the drivers of fire regimes in southern Russia, and particularly to identify the reasons for the marked increase in annual burned area since the mid-1990s (Dubinin et al., 2010).

Wildfire patterns are determined by many factors, but ultimately the key factors are fuel availability (moisture and amount) and ignition sources (Bond and van Wilgen, 1996; Meyn et al., 2007). In southern Russia and elsewhere, almost all ignitions are human-caused, and constant in number since human populations have remained fairly stable. Fuel availability, on the other hand, has

changed considerably over time, both in terms of fuel amounts and flammability, which in turn is determined by moisture. While fuel moisture is largely controlled by climate, the amount of fuel in grasslands is highly dependent on land use and especially grazing. Grazing influences the rate of vegetation recovery and directly reduces both amount and contiguity of fuels by consumption, wallowing, and trampling of vegetation, eventually leading to catastrophic shifts in vegetation structure and composition if grazing levels are not moderated (van de Koppel et al., 1997).

Most grasslands are experiencing intensifying land use, and long-term decrease in land use intensity is generally a rare phenomenon. However, periods of massive socioeconomic change may provide unique natural experiments (Diamond, 2001) to study the effects of decreasing land use intensity (e.g., lower grazing pressure) on fire patterns. Institutional changes in the former USSR have repeatedly led to changes in the vegetation, first after October Revolution and the Civil War of 1917-1920 and then after the end of World War II (1941-1945, Zonn, 1995b). Vegetation recovered after these events due to substantial decreases in human populations and livestock numbers, as well as the cessation of other agricultural activities. Vegetation changes were particularly evident in the arid grasslands of the Caspian plains because of the fragility of upper soil layer (largely alluvial sands).

The collapse of the former USSR in 1990/1991 also resulted in a sharp decline in livestock numbers, and detailed livestock data provided a much clearer picture of the changes in livestock numbers and hence grazing pressure for the 1990s than for prior periods of livestock declines (Figure 11). Soviet era livestock numbers were so high that they caused disastrous environmental effects (Vinogradov, 1995; Zonn, 1995a). Rangeland scientists had calculated that even on the best pastures, sheep density should not exceed 28-30 units/ha, but sheep density reached 130 heads/ha by 1975, and continued to increase until late 1980s (Zonn, 1995b). Overgrazing proved to be ecologically disastrous for the fragile arid ecosystems and ultimately led to desertification starting in

the 1960s and continuing until the early 1990s (Saiko and Zonn, 1997). The subsequent sharp decline in livestock numbers was paralleled by other substantial changes in ecosystem processes, including fire. During the period of overgrazing, fires were virtually absent in our study area (Dubinin et al., 2010). However, within half a decade after the declines in livestock numbers, annual burned area increased very rapidly (Figure 11), especially on sites that had previously been highly degraded (Dubinin et al., 2010).

The relationship between climate and burned area can be both direct and indirect. In seasonally dry, biomass-poor grasslands fuel amounts likely are more important drivers of fire than climate because fuels are desiccated during dry period, regardless of climate during the year (Meyn et al., 2007). However, the relative importance of climate may increase as land use intensity weakens (Meyn et al., 2007). Higher precipitation, especially during fire season can impede burning due to higher fuel moisture. On the other hand, high precipitation during the growing season prior to the fire season can lead to an increase in area burned because the rain increases fuel abundance (Flannigan and Wotton, 2001; Pausas, 2004). In other words, the amount of burning in a given year might depend on the weather during the previous year, which may determine fuel availability (Knapp 1998).

When examining the effects of climate on annual burned area it is important to consider global climate oscillation patterns as important explanatory factors (Flannigan and Wotton, 2001). Climate oscillations can be synchronized with fire weather conditions across large areas via teleconnection mechanisms. For example, El Niño - South Oscillation (ENSO) and North Atlantic Oscillation events are associated with increased fire activity in different regions of the world (Le Page et al., 2008; Greenville et al., 2009). Similar linkages occur between fire activity and the Arctic Oscillation (Balzter et al., 2005). In semi-arid and arid ecosystems, where precipitation is the limiting factor for fuel production and hence fires, increased rainfall might



result in a pulse of productivity and thus higher fuel availability in subsequent years (Holmgren et al., 2006). The relationship between broad-scale climate oscillations and local temperature and precipitation are regionally specific and additional analysis is required to establish how broad-scale climatic patterns affect fire patterns (Schonher and Nicholson, 1989; Keeley, 2004).

However, interactions between vegetation, climate, and burned area may be overridden by land use. Grazing pressure above a critical threshold can limit vegetation to such an extent that fires can no longer occur, irrespective of climatic conditions (van de Koppel et al., 1997). This means that climate probably would not predict burned area while livestock numbers were high under Soviet agricultural practices. However, once livestock numbers declined below a critical threshold vegetation recovered to a point where fires could be sustained, and it was after these thresholds had passed that climate would exert its effects on the burned area of a given year.

The goal of this study was to understand what explains fire regimes during rapidly changing socio-economic settings. Our general hypothesis was that socio-economic changes that followed the breakdown of the Soviet Union lowered grazing pressure, allowed vegetation to recover, and increased fuel availability, thereby providing the necessary prerequisite for fires, the amount of which was then controlled by climate. From this general hypothesis stemmed several specific hypotheses which we tested. We hypothesized that the burned area in a given year was:

- a) positively correlated with the amount of vegetation during the spring growing season immediately prior to the summer fire season, because of higher fuel availability;
- b) positively correlated with the amount of vegetation during the secondary growing season in the fall of the previous year, again due to higher fuel availability;
- c) positively correlated with climatic conditions that are favorable for vegetation growth during spring and fall of the previous year, also due to higher fuel availability;

d) more strongly correlated with climate during the period after livestock numbers had declined and grazing pressure lessened, because vegetation was able to recover to a point above which fires could occur;

e) negatively correlated with livestock numbers, because fewer livestock result in more abundant fuel;

and

f) negatively correlated with higher precipitation and lower temperature during fire season because higher fuel moisture impedes fires.

To test these hypotheses, we conducted statistical analyses to identify the best set of variables to explain annual burned area from 1985 to 2006. In addition, we conducted our analysis separately for the two time periods before and after the breakdown of the Soviet Union to test if the relationships between burned area and livestock, vegetation, and climate were changing from the Soviet to the post-Soviet period. To our knowledge, no previous studies of comparable temporal extent have attempted to explain burning dynamics in the grasslands of southern Russia and Central Asia.

## ***Methods***

### **Study area**

Our study area was located in the grasslands of Southern European Russia and occupied about 19,000 km<sup>2</sup> of the Republic of Kalmykia and Astrakhan Region (Figure 12). We had estimated annual burned area for this study area in previous research (Figure 11, Dubinin et al., 2010).

The climate of the study area is arid, with hot, dry summers (mean daily temperature of +24°C in July; max +44°C, 280 days of sunshine per year on average). Annual precipitation is 150 to 350 mm (with a mean of 286 mm from 1985 to 2007). Summer droughts are common, and most of the precipitation falls in spring and fall, coinciding with the two major growing seasons (Walter and Box, 1983). The topography is predominantly flat with a mean elevation of -15 m below sea level. The study area has a complex geological history of transgressions and regressions of the Caspian Sea. Soils are characterized by a gradient from sandy aeolian deposits and sandy loams in the southeast corner of the study area, to clay loam in the northwest (Kroonenberg et al., 1997).

Vegetation associations are typical for the northern Precaspian Plain and include both steppe and desert types. The main vegetation associations are shortgrass steppe (*Stipa* spp., *Festuca* spp., *Argopyron* spp., *Anizantha tectorum*, and other graminoids) and sage scrub (*Artemisia* spp., *Kochia prostrata*) (Golub, 1994). Shortgrass steppe is characterized by a short growing season in April and May and rapid senescence in the dry summer. The perennial grasses are well adapted to fire due to dense bunches, which protect seeds and basal meristems with tightly packed leaf-sheaths, and which can resprout after fire. Both annual and perennial grasses generate abundant fuels. Sagebrush (*Artemisia* spp.) dominated shrublands have less biomass, but a longer growing season, and sometimes exhibit a second vegetation peak in the fall and early winter (Kurinova and Belousova, 1989). *Artemisia* spp. is more susceptible to fire because its buds are situated above ground and can be killed or damaged by fires. The relative lack of fire tolerance by *Artemisia* spp. can lead to its substitution by *Stipa* spp. and other graminoids (Neronov, 1998). The primary human land use of the grasslands in the study area is for grazing

by domestic livestock, mainly sheep, and to a lesser extent cows and goats. The study area is sparsely populated (population density 0.8 to 1.4 persons/km<sup>2</sup>, (CIESIN and CIAT, 2005).

Broad-scale burning in the region can be traced as far back as 18<sup>th</sup> century (Pallas and Blagdon, 1802). However, little is known about spatial and temporal dynamics of fires both historically and in the recent past since practically no research has been done on fire patterns in southern Russia prior to our research and governmental statistics are largely inaccessible, incomplete, and biased (Dubinin et al., 2010).

#### *Study period and data sets used*

We studied burned area and its drivers from 1985 to 2006. The choice of the study period was determined by satellite image availability, and by the significant changes in land use that occurred after 1991. Our 22-year burned area record is one of longest derived from coarse resolution satellite data. Other burned area studies of similar long time spans have either focused on forests and mesic grasslands or used indirect or approximate estimates of burning (Beckage et al., 2003; Mouillot and Field, 2005; Chuvieco et al., 2008).

#### *Fire*

Remote sensing data are the primary source of information on burned area because they can provide both long and dense time series that can be used for statistical analysis (Kasischke and Penner, 2004). Historical and official records of burning in our study area are scarce and unreliable (Dubinin et al., 2010). Charcoal accumulation and tree-ring methods are not representative in grasslands because of low temporal resolution and low woody biomass. Our data set of burned area in Southern Russia consisted of 22 raster maps representing burned and non-burned areas for each year from 1985 to 2006. Burned area maps were generated based on

National Oceanic and Atmospheric Administration (NOAA) Advanced Very High Resolution Radiometer (AVHRR) Local Area Coverage data (Dubinin et al., 2010). The resolution of the burned area maps was 1.1 km pixel size. We mapped burned areas with supervised decision trees applied to an annual image pair representing one image taken during the peak growing season in the spring right before the fire season, and a second image taken right after the fire season in late summer (Dubinin et al., 2010).

### *Livestock*

Livestock population data originated from Ministry of Agriculture of the Republic of Kalmykia and represented the total population of sheep and goats for the study area. Though goats are combined with sheep in official statistics, they are rare (less than 4%). Cattle numbers are an order of magnitude smaller than those of sheep (Kalmstat, 2008).

### *Vegetation*

In terms of vegetation data, we focused on vegetation indices that served as a proxy for fuel amounts. The Normalized Difference Vegetation Index (NDVI) is significantly correlated with aboveground net primary production (ANPP) in many ecosystems, including grasslands (Tucker et al., 1985; Paruelo et al., 1997). We thus summarized vegetation as the mean monthly NDVI averaged for the entire study area. NDVI data were extracted from the GIMMS NDVI dataset (Tucker et al., 2005). GIMMS NDVI data were available twice a month from 1981 to 2006 and had 8-km resolution. GIMMS data are well suited for vegetation studies in semi-arid regions (Fensholt et al., 2009).

### *Climate*

Climate data were extracted from the Climate Research Unit Time Series product version 3 (Mitchell and Jones, 2005) which included 0.5° gridded mean (T) and maximum temperatures

(TMX), as well as precipitation data (PRE) interpolated from meteorological stations (New et al., 1999). In addition to local climate measures, we also looked at broad-scale climatic oscillation indices. Because it was not clear which broad-scale climatic pattern might affect local climate in our study area we chose three: the North Atlantic, the East Atlantic/Western Russia, and the Arctic oscillations index, abbreviated NAO, WR, AO respectively (Barnston and Livezey, 1987; Hurrell et al., 2003). The positive phase of AO indicates higher pressure in mid-latitudes, which results in more precipitation and enhanced greening in Central Asia during spring (Buermann et al., 2003). The positive phase of the NAO is associated with anomalous low pressure in the Subarctic, higher precipitation in the oceans between Scandinavia and Iceland and lower precipitation in Southern Europe (Hurrell et al., 2003). During positive phase of the WR, temperatures above North Caspian region and West Russia are drier than normal (Krichak et al., 2002). Climatic indices were averaged by season and year, thus each oscillation was represented by 5 variables. Additionally, we calculated seasonal and yearly precipitation and temperature summaries.

## **Statistical approach**

### *Bivariate analyses*

We evaluated the relationships between burned area and potential predictor variables with Pearson's correlations and tested our hypotheses with linear regression models with and without interaction terms. Log-transformed annual burned area was the response variable. Predictor variables included livestock, vegetation, and climate, both for the current and the previous year. Residuals were examined for possible temporal autocorrelation using Durbin-Watson tests and autocorrelative function plots (maximum time lag = 10 years). Heteroscedasticity was evaluated via visual

examination of normality plots. To identify the best predictors in our candidate models, we used best subsets analysis. The number of times that a particular variable entered the model was calculated for 20 best models. Models were ranked according to their adjusted r-squared values. All statistical analysis were performed in CRAN R (R Development Core Team, 2009).

Initial analyses showed that spring NDVI was a strong predictor of burned area, and this raised the question about what determined spring NDVI. We used the same statistical approach as outlined above, but used spring NDVI as the response variable, which was then modeled with the same set of explanatory factors, except for measures made later than spring.

Finally, we studied the effect of climate oscillations on local climate. Initial analyses showed strong correlations between climate oscillations at broad scales and local measures of temperature and precipitation.

#### *Regression modeling for before and after periods*

One of our hypotheses was that the relationship between climate and fire was fundamentally different before and after the breakdown of the Soviet Union, because the resulting declines in livestock numbers allowed vegetation to recover to a point above which fires were possible, and climate could exert control on burned area totals. To test this hypothesis, we stratified the data into two periods of time representing 'before' and 'after' the breakdown of the Soviet Union. We chose 1996 as a splitting year, resulting in 11 observations for the 'before' period and 12 observations for 'after'. We chose 1996 because previous research found that 5- to 6 years were required for the recovery of vegetation after grazing declined (Bananova, 1992; Zonn, 1995b). To assess how sensitive our results were in regards to the splitting year, , we calculated mean and standard deviation for a set of correlations separately for 'before' and 'after' periods while the splitting year was changed in annual steps from 1992 to 1997. To determine if slopes of the regressions

were significantly different between before and after periods we used regression models in the form of:

$$Y = I + x + dummy + dummy:x$$

Where  $x$  was a given independent variable, *dummy* indicated the period (0: before, and 1: after), and *dummy:x* represented an interaction between the dummy variable and the independent variable. We checked for changing variance structure by using a mixed-effect model that included group-effects. We used ANOVA to check if the more complex model was significantly different from the more simple one (i.e., the null hypothesis was that variance was not significantly different before and after). If we were unable to reject the null hypothesis, then we used simpler model, and otherwise the grouped variance mixed effect model. We were particularly interested to test if there were significantly different slopes in the regression lines for particular variables in the ‘before’ and ‘after’ periods. Only variables which showed statistically significant correlations in either of the periods were included in these analyses.

We also selected the best 2-variable regression model to estimate coefficients for two separate regressions, one for ‘before’ and one for ‘after’. We tested if coefficients were significantly different using Chow tests, commonly used in time series analysis to test for the presence of a structural break (Chow, 1960).

#### *Structural equation modeling*

Our general hypothesis was that socio-economic change, grazing pressure, vegetation change, fuel amounts, and climate all were related, and that their interactions ultimately determined burned area. The challenge was to identify these complex relationships. Regression analysis is well suited to identify significant predictors of burned area, but less powerful to assess the relationships and interactions among the predictor variables (and strong relationship raise collinearity concerns).



Structural equation modeling can examine possible causal pathways among intercorrelated variables, identify associations among variables while statistically controlling for other model variables (i.e., to partition relationships), and examine the likelihood of alternative models given the data at hand (Bollen, 1989). To facilitate our analyses and interpretations, we developed a meta-model, which represented main linkages between system components without statistical details. Our bivariate correlation analyses functioned as an indicator selection analysis for the structural equation model, and helped identify representatives for each entity of the conceptual model, which included global climate that influenced local climate. Local climate, livestock and burning all influenced fuels, and fuels in turn influenced burning. Significant variables indicated causal relationships and were used to construct a structural equation model (Grace, 2006). Nested models included a common set of variables and assumptions about dependencies (e.g., NDVI depends on livestock), but differed in the pathways deemed to be significant. We used maximum likelihood procedures to estimate coefficients and to evaluate model goodness-of-fit. Sequential application of single-degree-of-freedom  $\chi^2$  tests was used to determine which pathways should be retained in the final model. Decisions to keep or remove a variable or path from the model were also informed by modification indices (Jöreskog and Sörbom, 1984; Grace and Bollen, 2006). We conducted structural equation modeling in Amos 5.0 (Amos Development Corp, 2003).

## ***Results***

Our prior analysis of remote sensing data had shown that burned area increased very rapidly in southern Russia after 1996 (Figure 11, Dubinin et al, 2010). Statistical tests confirmed that burned area in the 1997-2006 period was significantly higher than in the 1985-1996 period even though there was a high variability among years ( $1044 \text{ km}^2 (\pm 1032)$  versus  $37 \text{ km}^2 (\pm 66)$ ,  $t =$

3.08,  $P = 0.013$ , Figure 13). The comparison of correlation coefficients with 1996 as a splitting year versus the mean and standard deviation of a set of correlations estimated while the splitting year was shifted from 1992 to 1997 confirmed that the use of 1996 as the splitting year did not bias our results (Table 3).

## **Livestock and fire**

Livestock numbers were significantly lower in the 1997-2006 period, and showed the strongest percent change (Figure 14). Both current and previous year livestock were negatively correlated with burned area, but correlations were only significant when analyzed for the entire study period and not significant when the dataset was split into ‘before’ and ‘after’ periods. Burned area was more strongly correlated with livestock numbers in the previous year than livestock numbers in the current year. Both current and previous year livestock were among the five top bi-variate models and statistically significant at the  $p < 0.05$  level (Table 4).

## **Vegetation and fire**

Mean April NDVI was significantly higher in the ‘after’ period compared to the ‘before’ period, but while May NDVI showed the same trend, the difference was statistically not significant (Figure 14 and Table 3). However, April NDVI and even more so May NDVI were the most powerful explanatory variables predicting burned area (May NDVI,  $p=0.0012$ ,  $r=0.66$ ) (Table 3). Correlation was particularly high for the ‘after’ period. However, neither spring NDVI measure was significantly correlated with burning in the ‘before’ period only. Among our other vegetation measures, during the ‘before’ period, only September NDVI in the previous year was significantly correlated with burned area. Slopes of the relationships between May NDVI and

fire were significantly different according to our models (Table 7). Best subsets analysis showed that May NDVI was present in 12 out of the 20 best candidate models (Table 8).

## Climate and fire

Correlations of burned area with precipitation and temperature were generally weaker than with vegetation. However, we did find a significant positive correlation for precipitation in April (Table 3). The only significant correlation with temperature was for August, which was positive. Specifically, the daily temperature range in August was more strongly correlated with burning than maximum temperature itself. Moreover, mean temperature, temperature range in August, and precipitation showed statistically significant differences between the ‘before’ and ‘after’ after periods, although the significance of the difference in precipitation was marginal (Figure 14). Summary climate averages were highly correlated with monthly values (i.e., the correlation between summarized maximum summer temperatures and maximum temperature in August was 0.71). Correlations between burned area and climate variables were similar in the ‘before’ and ‘after’ periods, and correlations were not significant. Only summarized summer precipitation was found to be significantly negatively correlated with burning during period ‘after’. Climate measures from the previous year were not significantly correlated with burned area either.

The relationships of burned area with climate oscillation indices were complex. Several climate oscillation indices were significantly correlated with burning either ‘before’ or ‘after’, but only the previous year fall Western Russia oscillation index showed any evidence of significant change between the two periods (Table 3). However, out of 6 indices that were significantly correlated with burned area, 3 were previous year fall indices. Previous year

average fall WR index was a particularly strong predictor and was included in 7 of the 20 best models (Table 8).

### **Relationships among spring vegetation, climate, and livestock**

Vegetation productivity in spring was consistently positively correlated with spring precipitation and several fall and winter oscillation indices, and consistently negatively correlated with livestock numbers in the previous year (Table 6). Not surprisingly, NDVI in May was strongly correlated with NDVI in April. However, precipitation in the previous year was better correlated with spring NDVI during the ‘before’ period and the entire time series, and the correlation was statistically not significant during the ‘after’ period. We also found several strong correlations between spring NDVI and climate oscillation indices for the previous year, especially during the ‘after’ period (Table 6).

### **Local and broad-scale climate**

High correlations between burned area and climate oscillations did not correspond to strong relationships between climate oscillation indices and local measures of temperature and precipitation. Though there were some statistically significant correlations between precipitation and temperature and oscillation indices for particular periods, correlations were weak (Table 5). For example, we found a significant negative correlation between both spring and summer maximum temperatures versus the NAO index in April, and cumulative summer precipitation was significantly correlated with the August WR index. However, none of the fall climate oscillation indices, which were well correlated with burned area, were significantly correlated with any precipitation or temperature measures. At the same time, there were a number of other correlations that were hard to interpret. For example, the April NO index was significantly

correlated with August temperatures, and the winter WR index was correlated with April precipitation. Spatial correlation fields calculated for fall precipitation and temperature and respective oscillation indices confirmed that correlation between broad scale climate and local climatic patterns in the study area were not statistically significant (Figure 15).

### **Multiple regressions**

The model yielding the highest adjusted  $r^2=0.73$  included previous year AO, NDVI5 and their interaction. However, previous year AO was not statistically significant by itself ( $p = 0.1$ ) and the interaction term was only marginally significant ( $p = 0.05$ ) at the 0.05 level. The model where all three parameters were statistically significant at 0.05 level was ranked third had adjusted  $r^2 = 0.58$  and included May NDVI, livestock population and their interaction (Table 2). Five best models with broad-scale climate indices omitted yielded smaller adjusted  $r^2$  values and were also dominated by NDVI.

### **Structural equation modeling**

The final structural equation model included 6 exogenous variables and one endogenous variable ( $\chi^2 = 7.7$ ,  $df = 13$ , Probability level = 0.862, Figure 16). Results provided by structural equation modeling clarified a number of relationships that were not clear from the univariate and multivariate analyses. The structural equation modeling suggested that livestock mainly influenced vegetation productivity in April, but not in May. Vegetation productivity in April in turn was more influenced by the cumulative precipitation of the previous year than by precipitation in April itself. Structural equation modeling confirmed that vegetation productivity

in May was dependent on April precipitation and on April NDVI. There was no indication that vegetation productivity in April by itself influenced burning. In models that included pathways between livestock and burned area, and between April NDVI and burned area, these pathways were all non-significant. The pathway between precipitation during the fire season and burned area was not significant either and we removed this variable. Because the bivariate analysis did not reveal any mechanisms by which climate oscillations influence burned area, we did not include oscillation indices in the structural equation modeling.

## ***Discussion***

Our study area experienced a rapid increase in burning following the collapse of the USSR, and our results highlight that these fire patterns resulted from interactions between decreasing livestock densities and recovering vegetation, modulated by climate. The structural equation models highlighted the complexity of the system, where the factors that influenced burning the most were dependent on a host of other factors.

## **Livestock**

Livestock was the main determinant of the amount of vegetation, as measured by NDVI. Our analysis supported the hypothesis that livestock numbers were significantly related to the increase in burned area through the amount of fuels. Though this relationship appears logical, the decrease of livestock and following increase in burning had rarely been documented explicitly and evidence is mostly historical (Savage and Swetnam, 1990). During the ‘before’ period high livestock numbers masked out the relationships between burned area and other variables such as climate. In contrast during the ‘after’ period, the release of livestock pressure has led to

accumulation of fuels and consequent increases in burning. This is different from what was observed in other areas of the world, for example in the Sahel, where lands remained barren for 20 years after overgrazing (Sinclair and Fryxell, 1985).

However, the correlation between burned area and livestock numbers was not as strong as we expected, possibly due to threshold effect (Turner and Gardner, 1991; van de Koppel et al., 1997). The pattern of the relationship suggests the existence of a critical threshold in livestock numbers that may have been reached in early 1990s: livestock density was no longer high enough to impact vegetation on a broad scale. The recovery of the vegetation was also coupled with successes in restoration activities and creation of nature reserve occupying a portion of the study area (Zonn, 1995b).

## Vegetation

The decrease in livestock numbers was accompanied by an increase in vegetation productivity, especially in spring. A strong relationship between spring NDVI and burning supported the interpretation that fuel amount before the fire season was significantly lower when livestock numbers were high and fire was not able to spread as far. However, while AVHRR NDVI was shown to be strongly related to primary productivity it is not a perfect measure of biomass (Box et al., 1989; Paruelo et al., 1997). The fact that we did not observe strong change in NDVI between 'before' and 'after' suggest that changes in composition and structure might have occurred. This is consistent with long term plot-level studies of grass-dominated communities after release of livestock pressure that reported increasingly higher presence of perennial fire adapted grasses, such as *Stipa capillata* (Neronov, 1998). Such change has resulted in considerable increase in biomass, from 0.5 to around 1 ton/ha aboveground biomass of *Stipa*

spp. in *Stipa* spp. dominated communities (Dzhapova, 2007). Increase in grassy fuels concomitant with an increase in burning is consistent with other parts of the world and represents a positive feedback loop (D'Antonio and Vitousek, 1992; Brooks et al., 2004).

Statistical evidence confirmed the hypothesis that burning was positively correlated with vegetation conditions during fall vegetation season of previous year. Burning during the 'before' period was linked to previous year vegetation productivity of the secondary growing period (i.e., fall NDVI). Fall NDVI represents secondary growth that is not consumed by fire and which becomes fuel in the form of the dry litter in the subsequent year. The lack of a relationship in the 'after' period can be explained by the increase in burning itself, leaving less vegetation for the secondary growing season. Alternatively, we could have missed some of the previous year effects due to litter accumulation, because non-photosynthetically active vegetation is not captured by NDVI but can contribute greatly to fire spread.

## **Climate**

Climate influenced burning both during the summer fire season and during spring green-up. During the fire season high precipitation and low temperatures led to increased fuel moisture and inhibited burning in some years (e.g., in 2003). Strong relationship existed between late summer maximum temperature and burning overall. However, summer precipitation was significantly correlated only during period 'after'. Comparison of change in climatic variables showed that the period 'after' experienced overall hotter summers.

High precipitation during green-up may have caused more fire due to more abundant fuel (Buermann et al., 2003). The amount of biomass in high precipitation years exceeds the average by 20-30% depending on vegetation type (Dzhapova, 2007). According to our correlation



analyses we found support for our hypothesis that spring precipitation was significantly related to burning. However, prior-year precipitation was only marginally ( $r = 0.42$ ) related to vegetation productivity in April. The lack of a relationship with prior-year precipitation suggests that after the decrease in livestock numbers, the amount of fuel increased to a point when fires occur in any year irrespective of additional precipitation during the previous year.

The representation of climate by station-based local climate measures could be unsatisfactory due to lack of stations and might be improved by using broad-scale climatic indices instead. Relationship between broad-scale climatic variables, such as previous year fall Western Russia oscillation, and burned area yielded significant correlations. Low values of WR index indicate milder and wetter weather in the region, which in turn might have lead both to more fuel production and increased fuel moisture during that year. The importance of both indices during ‘before’ period showed that fall precipitation was a more important determinant of burning before 1996. Secondary growth during fall period of the previous year became dry litter to burn the next year. The relationship between fall NDVI and climate variables is also supported by the relatively high correlation between previous year fall NDVI and burning during ‘before’ period. However, though such explanations were plausible, we were not able to confirm them with our precipitation data. Different oscillation indices often showed high correlations without clear linkages to local climate. High correlation was often found for seasonal indices that were timed well before or after corresponding measures of local climate. We thus caution that broad-scale climate variables should be used with care. In our case study, the mechanism of broad-scale climate influence on burning remained unknown, and this raised the question of how such indices were related to local climate in other studies, where the relationship with local climate

variables was not explored, but oscillation indices were used to explain burning (Greenville et al., 2009).

Overall the effect of current climate was stronger than that of climate in the previous year. This was not consistent with the findings from analogous ecosystems in the Great Plains of the U.S., California chaparral, and arid spinifex grasslands of Australia where one- and two-years previous year climate variables were better correlated with burning than current-year variables (Knapp, 1998; Keeley, 2004; Greenville et al., 2009). Our analysis of the relationship between climate and vegetation productivity showed that previous year climate had little effect on vegetation during the period of intensive burning. However, correlations of burned area with a range of broad-scale climatic indices, especially during fall, and the results from the structural equation modeling suggested that both precipitation and temperature data might not be adequate for our study area, or that the effect of broad-scale changes on fires is transmitted through other parameters that we did not measure, such as evapotranspiration.

In summary, we conclude that though climate was an important factor affecting burned area, especially in the Post-Soviet period, it could not explain the drastic increase in burned area between the two periods by itself. Substantial change in burning regime was driven by change in vegetation, which recovered as a consequence of relief from livestock pressure. If the trend of rebounding livestock numbers since 2000 continues in the future, then we may also expect a corresponding decline in burning in near future. In general, our results suggests that change in socio-economic conditions that lead to lower grazing pressure will likely lead to an increase in burning regardless of climate, and such declines in livestock grazing can be expected in particular after strong institutional changes.

## ***Acknowledgements***

We gratefully acknowledge support for this research by the NASA Land-Cover and Land-Use Change Program, a NASA Earth System Science Dissertation Fellowship for M. Dubinin (07-Earth07F-0053), and the National Geographic Society. We are thankful for J. Burton's help with structural equation modeling and N. Keuler assistance with statistics.

## ***Bibliography***

- Archer, S., Schimel, D.S., Holland, E.A., 1995. Mechanisms of shrubland expansion - land-use, climate or CO<sub>2</sub>. *Climatic Change* 29, 91-99.
- Archibald, S., Bond, W.J., 2004. Grazer movements: spatial and temporal responses to burning in a tall-grass African savanna. *International Journal of Wildland Fire* 13, 377-385.
- Balzter, H., Gerard, F.F., George, C.T., Rowland, C.S., Jupp, T.E., McCallum, I., Shvidenko, A., Nilsson, S., Sukhinin, A., Onuchin, A., Schmulius, C., 2005. Impact of the Arctic Oscillation pattern on interannual forest fire variability in Central Siberia. *Geophysical Research Letters* 32.
- Bananova, V.A., 1992. Antropogenic desertification of arid grasslands of Kalmykia. Doctoral thesis. Institute of deserts, Ashkhabad.
- Barnston, A.G., Livezey, R.E., 1987. Classification, Seasonality and Persistence of Low-Frequency Atmospheric Circulation Patterns. *Monthly Weather Review* 115, 1083-1126.
- Beckage, B., Platt, W.J., Slocum, M.G., Pank, B., 2003. Influence of the El Nino Southern Oscillation on fire regimes in the Florida everglades. *Ecology* 84, 3124-3130.
- Bollen, K.A., 1989. Structural equations with latent variables. John Wiley and Sons, New York.

- Bond, W.J., Keeley, J.E., 2005. Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems. *Trends in Ecology & Evolution* 20, 387-394.
- Bond, W.J., van Wilgen, B.W., 1996. *Fire and plants*. Chapman and Hall, London.
- Box, E.O., Holben, B.N., Kalb, V., 1989. Accuracy of the AVHRR vegetation Index as a predictor of biomass, primary productivity and net Co<sub>2</sub> flux. *Vegetatio* 80, 71-89.
- Brooks, M.L., D'Antonio, C.M., Richardson, D.M., Grace, J.B., Keeley, J.E., DiTomaso, J.M., Hobbs, R.J., Pellant, M., Pyke, D., 2004. Effects of invasive alien plants on fire regimes. *Bioscience* 54, 677-688.
- Buermann, W., Anderson, B., Tucker, C.J., Dickinson, R.E., Lucht, W., Potter, C.S., Myneni, R.B., 2003. Interannual covariability in Northern Hemisphere air temperatures and greenness associated with El Nino-Southern Oscillation and the Arctic Oscillation. *Journal of Geophysical Research-Atmospheres* 108.
- Chow, G.C., 1960. Tests of Equality Between Sets of Coefficients in Two Linear Regressions. *Econometrica* 28, 591-605.
- Chuvieco, E., Englefield, P., Trishchenko, A.P., Luo, Y., 2008. Generation of long time series of burn area maps of the boreal forest from NOAA-AVHRR composite data. *Remote Sensing of Environment* 112, 2381-2396.
- CIESIN, CIAT, 2005. Gridded Population of the World Version 3 (GPWv3): Population Density Grids. Socioeconomic Data and Applications Center (SEDAC), Columbia University, Palisades, NY; Available at <http://sedac.ciesin.columbia.edu/gpw>. Last accessed: 06.28.2009.
- Csiszar, I., Denis, L., Giglio, L., Justice, C.O., Hewson, J., 2005. Global fire activity from two years of MODIS data. *International Journal of Wildland Fire* 14, 117-130.

- D'Antonio, C.M., Vitousek, P.M., 1992. Biological Invasions by Exotic Grasses, the Grass Fire Cycle, and Global Change. *Annual Review of Ecology and Systematics* 23, 63-87.
- Diamond, J., 2001. Ecology: Dammed Experiments! *Science* 294, 1847-1848.
- Dubinin, M., Potapov, P., Lushchekina, A., Radeloff, V.C., 2010. Reconstructing long time series of burned areas in arid grasslands of southern Russia by satellite remote sensing. *Remote Sensing of Environment*, doi:10.1016/j.rse.2010.02.010.
- Dzhapova, R.R., 2007. Dynamics of vegetative cover of Ergeni highlands and Caspian lowlands within Republic of Kalmykia. Moscow State University, Moscow.
- Fensholt, R., Rasmussen, K., Nielsen, T.T., Mbow, C., 2009. Evaluation of earth observation based long term vegetation trends - Intercomparing NDVI time series trend analysis consistency of Sahel from AVHRR GIMMS, Terra MODIS and SPOT VGT data. *Remote Sensing of Environment* 113, 1886-1898.
- Flannigan, M.D., Wotton, B.M., 2001. Climate, weather, and area burned. In: Johnson, E.A., Miyanishi, K. (Eds.), *Forest fires*. Academic Press, San Diego.
- Golub, V.B., 1994. The desert vegetation communities of the Lower Volga Valley. *Feddes Repertorium* 105, 499-515.
- Grace, J., 2006. *Structural equation modeling and natural systems*. Cambridge University Press, Cambridge, UK.
- Grace, J.B., Bollen, K.A., 2006. The interface between theory and data in structural equation models. Open-File Report 2006-1363. U.S. Geological Survey, Reston, Virginia.
- Greenville, A.C., Dickman, C.R., Wardle, G.M., Letnic, M., 2009. The fire history of an arid grassland: the influence of antecedent rainfall and ENSO. pp. 631-639.

- Guevara, J.C., Stasi, C.R., Wuilloud, C.F., Estevez, O.R., 1999. Effects of fire on rangeland vegetation in south-western Mendoza plains (Argentina): composition, frequency, biomass, productivity and carrying capacity. *Journal of Arid Environments* 41, 27-35.
- Holmgren, M., Stapp, P., Dickman, C.R., Gracia, C., Graham, S., Gutierrez, J.R., Hice, C., Jaksic, F., Kelt, D.A., Letnic, M., Lima, M., Lopez, B.C., Meserve, P.L., Milstead, W.B., Polis, G.A., Previtalli, M.A., Michael, R., Sabate, S., Squeo, F.A., 2006. Extreme climatic events shape arid and semiarid ecosystems. *Frontiers in Ecology and the Environment* 4, 87-95.
- Hurrell, J.W., Kushnir, Y., Ottersen, G., Visbeck, M., 2003. An Overview of the North Atlantic Oscillation. In: Hurrell, J.W., Kushnir, Y., Ottersen, G., Visbeck, M. (Eds.), *The North Atlantic Oscillation: Climate Significance and Environmental Impact*.
- Jöreskog, K.G., Sörbom, D., 1984. LISREL VI User's guide. University of Uppsala, Dept. of Statistics, Uppsala.
- Kalmstat, 2008. Livestock census of 2008. ROSSTAT, Elista.
- Kasischke, E.S., Penner, J.E., 2004. Improving global estimates of atmospheric emissions from biomass burning. *Journal of Geophysical Research-Atmospheres* 109.
- Keeley, J.E., 2004. Impact of antecedent climate on fire regimes in coastal California. *International Journal of Wildland Fire* 13, 173-182.
- Knapp, P.A., 1998. Spatio-temporal patterns of large grassland fires in the Intermountain West, USA. *Global Ecology and Biogeography* 7, 259-272.
- Krichak, S.O., Kishcha, P., Alpert, P., 2002. Decadal trends of main Eurasian oscillations and the Eastern Mediterranean precipitation. *Theoretical and Applied Climatology* 72, 209-220.
- Kroonenberg, S.B., Rusakov, G.V., Svitoch, A.A., 1997. The wandering of the Volga delta: A response to rapid Caspian sea-level change. *Sedimentary Geology* 107, 189-209.

- Kurinova, Z.S., Belousova, Z.N., 1989. Effective use of forage resources. Kalmyk book publishing, Elista.
- Le Page, Y., Pereira, J.M.C., Trigo, R., da Camara, C., Oom, D., Mota, B., 2008. Global fire activity patterns (1996-2006) and climatic influence: an analysis using the World Fire Atlas. *Atmospheric Chemistry and Physics* 8, 1911-1924.
- Letnic, M., 2004. Cattle grazing in a hummock grassland regenerating after fire: The short-term effects of cattle exclusion on vegetation in south-western Queensland. *Rangeland Journal* 26, 34-48.
- Meyn, A., White, P.S., Buhk, C., Jentsch, A., 2007. Environmental drivers of large, infrequent wildfires: the emerging conceptual model. *Progress in Physical Geography* 31, 287-312.
- Mitchell, T.D., Jones, P.D., 2005. An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *International Journal of Climatology* 25, 693-712.
- Mouillot, F., Field, C.B., 2005. Fire history and the global carbon budget: a 1 degrees x 1 degrees fire history reconstruction for the 20th century. *Global Change Biology* 11, 398-420.
- Neronov, V.V., 1998. Antropogenic grasslands encroachment of desert pastures of North-Western Precaspian plains. *Successes of Modern Biology* 118, 597-612 (in Russian).
- New, M., Hulme, M., Jones, P., 1999. Representing twentieth-century space-time climate variability. Part I: Development of a 1961-90 mean monthly terrestrial climatology. *Journal of Climate* 12, 829-856.
- Pallas, P.S., Blagdon, F.W., 1802. Travels through the southern provinces of the Russian Empire, in the years 1793 and 1794. Printed by A. Strahan ... for T.N. Longman and O. Rees ... London.

- Paruelo, J.M., Epstein, H.E., Lauenroth, W.K., Burke, I.C., 1997. ANPP estimates from NDVI for the Central Grassland Region of the United States. *Ecology* 78, 953-958.
- Pausas, J.G., 2004. Changes in fire and climate in the eastern Iberian Peninsula (Mediterranean basin). *Climatic Change* 63, 337-350.
- Pyne, S.J., 1984. *Introduction to Wildland Fire*. John Wiley and Sons, New York.
- R Development Core Team, 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. URL <http://www.R-project.org>.
- Saiko, T., Zonn, I., 1997. Europe's first desert. In: Glantz, M., Zonn, I. (Eds.), *Scientific, Environmental and Political Issues in the Circum-Caspian Region*. Kluwer Academic Publishers, Dordrecht, Boston and London, pp. 141-144.
- Savage, M., Swetnam, T.W., 1990. Early 19th-Century Fire Decline Following Sheep Pasturing in a Navajo Ponderosa Pine Forest. *Ecology* 71, 2374-2378.
- Schonher, T., Nicholson, S.E., 1989. The Relationship between California Rainfall and ENSO Events. pp. 1258-1269.
- Sinclair, A.R.E., Fryxell, J.M., 1985. The Sahel of Africa - Ecology of a Disaster. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 63, 987-994.
- Stohl, A., Berg, T., Burkhardt, J.F., Fjaeraa, A.M., Forster, C., Herber, A., Hov, O., Lunder, C., McMillan, W.W., Oltmans, S., Shiobara, M., Simpson, D., Solberg, S., Stebel, K., Strom, J., Torseth, K., Treffeisen, R., Virkkunen, K., Yttri, K.E., 2007. Arctic smoke - record high air pollution levels in the European Arctic due to agricultural fires in Eastern Europe in spring 2006. *Atmospheric Chemistry and Physics* 7, 511-534.
- Tansey, K., Gregoire, J.M., Stroppiana, D., Sousa, A., Silva, J., Pereira, J.M.C., Boschetti, L., Maggi, M., Brivio, P.A., Fraser, R., Flasse, S., Ershov, D., Binaghi, E., Graetz, D., Peduzzi, P.,



2004. Vegetation burning in the year 2000: Global burned area estimates from SPOT VEGETATION data. *Journal of Geophysical Research-Atmospheres* 109.
- Tucker, C.J., Pinzon, J.E., Brown, M.E., Slayback, D.A., Pak, E.W., Mahoney, R., Vermote, E.F., El Saleous, N., 2005. An extended AVHRR 8-km NDVI dataset compatible with MODIS and SPOT vegetation NDVI data. *International Journal of Remote Sensing* 26, 4485-4498.
- Tucker, C.J., Vanpraet, C.L., Sharman, M.J., Vanittersum, G., 1985. Satellite Remote-Sensing of Total Herbaceous Biomass Production in the Senegalese Sahel - 1980-1984. *Remote Sensing of Environment* 17, 233-249.
- Turner, M.G., Gardner, R.H., 1991. Quantitative methods in landscape ecology: an introduction. In: Turner, M.G., Gardner, R.H. (Eds.), *Quantitative methods in landscape ecology*. Springer-Verlag, New York.
- van de Koppel, J., Rietkerk, M., Weissing, F.J., 1997. Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends in Ecology & Evolution* 12, 352-356.
- van der Werf, G.R., Randerson, J.T., Giglio, L., Collatz, G.J., Kasibhatla, P.S., Arellano, A.F., 2006. Interannual variability in global biomass burning emissions from 1997 to 2004. *Atmospheric Chemistry and Physics* 6, 3423-3441.
- Vinogradov, B.V., 1995. Forecasting dynamics of deflated sands of Black Lands, Kalmykia using aerial photography. In: Zonn I.S., N., V.M. (Ed.), *Biota and Environment of Kalmykia*, Moscow, Elista (in Russian), pp. 259-268.
- Walter, H., Box, E., 1983. Overview of Eurasian continental deserts and semideserts. *Ecosystems of the World*. Elsevier, Amsterdam, pp. 3-269.
- Zonn, I.S., 1995a. Desertification in Russia: Problems and solutions (an example in the Republic of Kalmykia-Khalmg Tangch). pp. 347-363.

Zonn, S.V., 1995b. Desertification of nature resources in Kalmykia during last 70 years and protection measures., Biota and Environment of Kalmykia, Moscow, Elista (in Russian), pp. 12-52.

## Tables

Table 3. Pearson correlations between explanatory variables and burned area for the ‘before’ (1983-1996), and ‘after’ period (1997-2006), and the entire time series, and the stability (i.e., the mean and standard deviation) of the correlation coefficients when moving the split year (1992-1997). T-test shows p-value for significance of the differences between before and after datasets. Significance levels of the univariate correlations between the explanatory variables and burned area are denoted as: \*\* < 0.01, \* < 0.05. Notation and abbreviations: °- variable lagged one year back, WR – West Russia, AO – Arctic Oscillation, NAO – North Atlantic Oscillation, tmx – maximum temperature, pre – precipitation, NDVI – Normalized Difference Vegetation Index. Numbers indicate period or specific month.

variable	r-before	r-after	r-full	t-test	r-before stability	r-after stability
WRspring	0.13	-0.73*	-0.04	0.27	0.09±0.13	-0.40±0.32
WRwinter°	0.34	-0.68*	-0.18	0.57	0.26±0.09	-0.47±0.18
WRfall°	-0.67*	-0.31	-0.62**	0.04	-0.79±0.07	-0.39±0.11
AOfall°	-0.43	0.83**	-0.05	0.11	-0.67±0.17	0.56±0.27
NAOfall°	-0.32	0.70*	-0.13	0.12	-0.48±0.12	0.50±0.21
tmx8	0.34	-0.09	0.48*	<0.01	0.39±0.06	0.18±0.22
pre4	0.38	0.42	0.57**	0.04	0.23±0.21	0.60±0.14
pre6-8	0.40	-0.65*	-0.21	0.99	0.52±0.08	-0.55±0.18
NDVI4	0.14	0.64*	0.55*	0.04	0.22±0.08	0.61±0.04
NDVI5	0.24	0.89**	0.66**	0.06	0.18±0.10	0.88±0.01
NDVI9°	0.63*	0.14	0.06	0.18	0.79±0.12	-0.06±0.16

livestock°	0.38	0.13	-0.52*	<0.01	0.06±0.22	-0.29±0.34
------------	------	------	--------	-------	-----------	------------

.

Table 4. Five best multivariate models explaining variability in log-transformed burned area a) with b) without broad climate variables (r denotes Pearson correlation between variable 1 and variable 2, significance levels: \*\* < 0.01, \* < 0.05).

a)

response	variable 1	variable 2	interaction	adj.r <sup>2</sup>	r
fire	AO <sup>°</sup>	NDVI5**	AO:NDVI5	0.73	0.23
fire	AOspring <sup>°</sup>	NDVI5**	AOspring <sup>°</sup> :NDVI5*	0.64	0.01
fire	livestock*	NDVI5**	livestock:NDVI5*	0.58	-0.37
fire	livestock <sup>°</sup> *	NDVI5*	livestock <sup>°</sup> :NDVI5*	0.58	-0.43
fire	AOspring <sup>°</sup>	pre4**	laospr:pre4*	0.57	0.14

b)

response	variable 1	variable 2	interaction	adj.r <sup>2</sup>	r
fire	livestock*	NDVI5**	livestock: NDVI5*	0.58	-0.37
fire	livestock*	NDVI5*	livestock: NDVI5*	0.58	-0.43
fire	tmx8 <sup>°</sup> *	NDVI5*	tmx8 <sup>°</sup> : NDVI5*	0.56	-0.00
fire	NDVI5*	NDVI6	NDVI5: NDVI6	0.50	0.52
fire	NDVI5	NDVI9	NDVI5:NDVI6	0.48	0.00

Table 5. Correlations between fire, local climate measures and oscillation indices AO, NAO, EAWR (monthly values, summarized by season: W – winter, Sp – spring, Su – summer, F - fall and yearly average), number indicates month (significance levels: \*\* < 0.01, \* < 0.05).

	tmx8	t3-6	t6-8	t9-11	t.sum	pre4	pre3-6	pre6-8	pre9-11	pre.sum
AO4	-0.20	0.05	-0.08	0.27	0.11	0.26	-0.17	0.26	-0.03	-0.01
AO8	-0.10	0.21	0.11	0.34	0.15	0.25	-0.30	-0.07	-0.22	-0.47
AO	0.15	0.17	-0.12	0.22	0.13	0.26	-0.37	0.43	0.10	0.17
AO										
winter	-0.17	0.12	-0.21	-0.04	0.09	0.17	0.11	0.21	0.21	0.09
AO										
spring	-0.26	0.08	-0.30	0.31	0.08	-0.06	-0.33	0.34	0.04	0.15
AO										
summer	0.07	0.24	0.10	<b>0.48*</b>	0.25	-0.27	-0.28	0.08	-0.23	-0.16
AOfall	0.13	-0.02	-0.18	-0.02	-0.01	-0.29	-0.01	0.16	-0.14	0.34
			-							
NO4	<b>-0.48*</b>	<b>-0.57*</b>	<b>0.63**</b>	0.06	-0.41	-0.01	-0.01	0.19	0.15	0.39
NO8	<b>-0.44*</b>	0.12	-0.16	0.27	0.18	0.14	-0.30	0.05	0.13	-0.42
NO	-0.21	-0.02	-0.36	0.02	0.01	-0.35	-0.17	0.08	0.04	0.21
NO										
winter	0.07	0.41	0.05	0.15	0.33	0.23	-0.11	0.26	-0.07	-0.11
NO										
spring	-0.42	-0.34	<b>-0.51*</b>	-0.17	-0.31	-0.30	-0.03	-0.13	0.15	0.19

NO						-				
summer	-0.18	0.20	-0.15	<b>0.61**</b>	0.28	<b>0.58**</b>	-0.33	0.25	0.06	-0.03
NOfall	0.04	0.11	-0.09	-0.24	0.13	-0.07	0.06	-0.11	-0.04	0.18
WR4	-0.24	<b>-0.51*</b>	-0.25	-0.34	-0.43	0.06	0.19	-0.10	0.36	0.03
WR8	-0.19	0.37	0.16	-0.05	0.21	-0.26	-0.14	<b>-0.46*</b>	-0.21	<b>-0.52*</b>
WR	-0.31	-0.24	<b>-0.47*</b>	-0.23	-0.40	-0.03	-0.14	-0.08	0.09	0.10
WR										
winter	-0.08	-0.36	-0.25	-0.24	-0.37	<b>0.54*</b>	-0.04	-0.02	0.10	0.18
WR										
spring	-0.14	-0.20	-0.26	-0.18	-0.07	-0.05	0.04	-0.05	<b>0.52*</b>	0.15
WR										
summer	-0.18	0.05	-0.30	0.00	-0.05	-0.15	-0.29	4	0.08	-0.07
WRfall	-0.27	0.00	-0.32	-0.29	-0.35	-0.18	0.05	-11	-0.28	0.07

---

Table 6. Relationships between a) April and b) May NDVI and other variables.

a)

	r-before	r-after	r-full
WRfall	-0.40	-0.68*	-0.62**
AOfall <sup>°</sup>	-0.45	0.79*	-0.05
NAOfall <sup>°</sup>	-0.26	0.69*	-0.13
pre4	0.54	-0.26	0.57*
pre.sum <sup>°</sup>	0.65*	0.39	0.42
livestock <sup>°</sup>	-0.53	0.04	-0.52*
NDVI10 <sup>°</sup>	0.64*	0.25	0.08

b)

	r-before	r-after	r-full
WRwinter <sup>°</sup>	0.41	-0.67*	-0.18
WRfall <sup>°</sup>	-0.17	-0.44	-0.62**
AOwinter <sup>°</sup>	0.62*	-0.27	-0.21
AOfall <sup>°</sup>	-0.16	0.91**	-0.05
AO <sup>°</sup>	0.66*	-0.04	-0.38
NAOfall <sup>°</sup>	-0.02	0.75**	-0.13
pre4	0.87**	0.15	0.57**
pre3-6	0.61	-0.06	0.14
pre.sum	0.41	-0.82**	-0.04
pre3-6 <sup>°</sup>	0.73**	0.23	0.17



pre.sum <sup>o</sup>	0.72**	0.58	0.42
livestock <sup>o</sup>	-0.23	0.05	-0.52**
NDVI4	0.71**	0.83**	0.54**

---

Table 7. Statistical significance of difference in slopes. Dummy – statistical significance for keeping 0/1 variable representing periods ‘before’ and ‘after’. Slopes – statistical significance for different slopes. Significance levels: \*\* < 0.01, \* < 0.05

	dummy slope		adj.r2
NDVI5		*	0.65
AOfall <sup>o</sup>	**	**	0.65
pre.sum	**	**	0.61
NDVI6 <sup>o</sup>	**	**	0.60
NAOfall <sup>o</sup>	**	*	0.56
pre6-8	**	*	0.56
WRwinter <sup>o</sup>	**	*	0.56
WRspring	**	*	0.56
NAO	**	*	0.51

Table 8. Results of two-variable best-subset analysis; only variables which participated in more than 1 model are shown.

variable	number of models
NDVI5	12
WRfall <sup>o</sup>	7
pre4	3
AOspring <sup>o</sup>	2
AO <sup>o</sup>	2

## Figures

Figure 11. Sheep population and burned area trends in the study area.




Figure 12. Study area boundary and location.



Figure 13. Relationships between selected explanatory variables and log-transformed burned area . Solid dots and solid line represent values and regression line for period ‘before’ (1986-1996), hollow circles and dashed line represent values for period ‘after’ (1997-2006).




Figure 14. Change in (a) burned area, (b) maximum August temperatures, (c) livestock numbers, and (d) April NDVI between ‘before’ and ‘after’ periods.






Figure 15. Correlation fields of broad-scale climatic oscillation patterns and local climate during fall months. Red indicates positive correlations, blue negative correlation, light colors - no statistical significance, and dark colors - statistical significance at 0.05 level, black – not analyzed.


a) AO – precipitation




b) NAO – precipitation




c) WR – precipitation



d) AO – temperature



e) NAO- temperature



f) WR- temperature








Figure 16. On top, (a) the final structural equation model, and below (b) full model with pathways and variables that were not retained (shown as dashed lines). Straight single-headed arrows represent significant effects of one variable on another ( $\alpha = 0.05$ ), while curved double-headed arrows represent correlations between variables. The relative strength of the effect is indicated by a path coefficient. Only pathways that are significant at  $\alpha = 0.05$  are included. See Table 1 for notations and abbreviations.

a)



b)



## **Chapter 3: Rapid vegetation change after land use changes and increasing wildfire activity in arid ecosystems of Southern Russia**

### ***Abstract***

Vegetation composition is an important characteristic of ecosystems and is strongly affected by disturbance regimes. In semi-deserts, the ratio of shrubs to grasses in particular is closely linked to disturbance especially wildfire. Shrub-encroachment is a global phenomenon in semi-deserts, however shrublands may also be replaced by grasslands in response to increasing fire activity and – potentially – changes in land use. The effect of fires on vegetation in semi-deserts is well understood, but effects of land use are less clear. In particular, it is unknown how sharp declines in land use intensity such as grazing, and subsequent increases in fire activity due to increasing fuel loads, affect vegetation communities, because sharp declines in land use intensity are rare. Our question here was if a sharp decrease in land use intensity (i.e., a 90% decline in sheep numbers in Kalmykia, southern Russia, following the collapse of the Soviet Union), and a subsequent increase in burning has resulted in a widespread shift from shrublands to grasslands and if this vegetation shift is related to legacies of past land use. Our study area was located in the Precaspian plain in the Republic of Kalmykia in southern Russia and encompassed 18,500 km<sup>2</sup>. We used satellite images, field data, and thematic maps to quantify change in vegetation communities, and to relate these changes to past and recent disturbances. Landsat TM satellite data for 1985, 1998, and 2007 were classified using Support Vector Machines and results were compared to the perimeters of burned areas and former open sands. We found a modest increase

in the area covered by grasslands from 1985 to 1998, and a substantial increase from 1998 to 2007. The shift from shrubs (*Artemisia* spp.) to grasses (*Stipa* spp.) was common in areas that had been plowed and then abandoned in the 1950s. However, these areas constituted less than 20 % and 10 % of the area that transitioned to grasslands by 1998 and 2007 respectively. Most of the cumulative burned areas of 1998-2006, were coincident with grasslands, especially in the southern part of our study area (on sandy soils). However, burned areas in the northern part did not convert to grasslands, and some areas in the southern part converted without evidence of fire. The observed vegetation changes in total represent considerable ecosystem changes in the Precaspian region of Russia. Decreasing grazing pressure, subsequent increases in burning, and long-term land use legacies resulted in a transition from shrub-dominated vegetation types to grass-dominated ones. Ultimately, these results highlight that the socio-economic changes that followed the collapse of the Soviet Union, resulting in strong declines in grazing pressure, caused a switch of ecosystem types due to the interaction of grazing, fire, and vegetation.

## ***Introduction***

In human-dominated landscapes, vegetation communities tend to homogenize and become less diverse if disturbance pressure is uniformly intensive (Olden et al. 2004). Particularly high disturbance pressure may induce ecosystem shifts (Scheffer et al. 2001; van de Koppel et al. 1997). However, it is often unclear if ecosystems can recover if disturbance pressure subsides. The declines in particular species that accompany ecosystem shifts may hamper an ecosystem's ability to recover and to provide valuable ecosystem services such as water, grazing capacity, and soil organic matter.

In semi-deserts, vegetation declines that follow increasing disturbance (such as grazing) are well documented, but studies that show if and how semi-deserts recover when disturbance pressure is released are less common. The ratio of shrubs to grasses projective cover is an important indicator of the state of the vegetation in semi-deserts and changes in this ratio over time can elucidate ecosystem trends, and reflect the effects of historic and recent disturbances. Broadly speaking, there are two trajectories semi-desert ecosystems can follow. First, semi-deserts can undergo shrub encroachment. Currently, shrub-encroachment is a global phenomenon, occurring, for example, in the grasslands of Australia, Africa, USA, and India (Archer et al. 1995; Roques et al. 2001; Van Auken 2000). The increase in woody biomass has been attributed to increased CO<sub>2</sub> concentrations and changes in rainfall patterns, but the most common causes are the suppression of fires, and/or increased land use intensity, particularly grazing, which allows shrubs to take over (Archer 1994; Schlesinger et al. 1990; Van Auken 2009).

The other broad trend is the replacement of shrublands by grasslands, which can be the result of frequent burning (Bond et al. 2005; Callaway and Davis 1993; Grigulis et al. 2005).

Increases in grass cover in the semi-deserts of the Western hemisphere due to fires are particularly striking in the case of invasive annual grasses, such as cheatgrass (*Bromus tectorum*), and cheatgrass spread is both a drivers and a result of increase in fire frequency (Brooks et al. 2004; HilleRisLambers et al. 2010).

The relationship of fire and vegetation structure and composition might also be influenced by past land use. Long-term land-use legacy effects are well document in forests (Foster et al. 1998; Zarin et al. 2005). Semi-deserts might be as sensitive to prior land use. Indeed, which species are most abundant in semi-deserts is strongly related to the recent grazing history (Milchunas and Lauenroth 1993). However, effects of prior land use are often masked by gradual vegetation changes.

Vegetation change in arid ecosystems is usually studied either using historical approaches (Archer 1994; Mack 1981), via conceptual models (van de Koppel et al. 1997; van Langevelde et al. 2003), or with field plots and especially exclosures, which allows tracking of changes explicitly (Belsky 1992; Milchunas and Lauenroth 1993). In regard to land use, the dominant question has been how increases in land use intensity, and especially in grazing, affect vegetation. Landscape-scale studies focusing explicitly on the response of ecosystems to decreases in land use intensity over large areas and long (20+ year) time spans are relatively rare. That is unfortunate, because sheep numbers have declined rapidly in the last two decades across the globe (Figure 17), and since sheep are the main livestock species in semi-deserts, this may have resulted in substantial vegetation changes. However, the results of the decline in sheep numbers are unclear, and hard to predict, because there may be lag effects in the response of fire and vegetation to changes in land use (Dubinin et al. in review).

Russia represents a unique opportunity to study the effect of both decreasing land use intensity and long-term land use legacies on fire and ultimately vegetation communities. While livestock numbers, specifically sheep, have decreased throughout the world, former Soviet Union countries represent the case where the decline has been particularly rapid: the Russian Federation and Kazakhstan occupy two out of three top spots among greatest ‘losers’ of sheep, and they lost most of their sheep in less than a decade (Figure 17). It is unknown how this decline translated into ecosystem changes, especially vegetation. The Republic of Kalmykia, located in the south of Russia is particularly interesting with regard to changes in livestock numbers because it is the biggest livestock producer together with the neighboring Republic of Dagestan (ROSSTAT 2006). In Kalmykia, environmental mismanagement started in 1960s, when increasing livestock numbers could not be supported by natural pastures alone, and substantial amounts of lands were plowed to grow forage crops (Zonn 1995a). Ultimately, these efforts failed, large areas became void of all vegetation, and forage production ceased almost completely by the 1980s. The land use legacy of the areas used in the past for forage crops is unknown though.

Similarly, it is unknown how decreasing land use intensity in Kalmykia affected the vegetation. Sheep density was as high as 130 head/ha in the 1970s and 1980s, but livestock numbers declined sharply after the collapse of the former USSR in 1990/1991, dropping from 3.2 to 0.6 million heads in 10 years (Zonn 1995b). The decline in grazing pressure resulted in higher vegetation productivity as measured by NDVI (Dubinin et al. 2010a), and also a very strong increase in the area burned each year (Dubinin et al. 2010b). There were very few fires before 1997, but the area burned annually increased almost exponentially thereafter, albeit with high variability in burned area from year to year (Dubinin et al. 2010b). Field studies suggested that there were an increase in grass cover, especially *Stipa* spp., as well as *Bromus tectorum*

(Neronov 1998; Shilova et al. 2007). However, prior studies of recent vegetation change in Kalmykia were small in geographic extent (Hoelzel et al. 2002; Neronov 1998), and provide an incomplete spatial picture of the vegetation changes that occurred.

Our overarching goal was to understand the effect of change in land use on semi-desert vegetation communities. Our specific objectives were: 1) to quantify change in different vegetation types for two periods: from 1985 to 1998 and from 1998 to 2007; 2) to relate the vegetation changes to spatial patterns of burning and to land use legacies, especially past attempts of row crop agriculture. Given the substantial declines in livestock and the subsequent increase in burning we hypothesized that: 1) the area dominated by grasses increased and the area dominated by shrubs declined; 2) the area of bare, open sands decreased; and 3) areas where grasses replaced shrubs were associated with both wildfires and formerly open sands.

## ***Methods***

### **Study area**

Our study area was located in southern European Russia and occupied about 31,485 km<sup>2</sup> of the Republic of Kalmykia and Astrakhan Region (Figure 18). The study area is sparsely populated (population density 0.8 to 1.4 persons/km<sup>2</sup> (CIESIN and CIAT 2005). Contrary to expected worldwide trends in population growth in arid and semi-arid regions (Warren et al. 1996) no substantial population growth is expected in our study area (CIESIN et al. 2005).

The climate of the study area is arid, with hot, dry summers (mean daily temperature of +24°C in July; max +44°C, 280 clear days per year on average). Annual precipitation is 150 to 350 mm (with a mean of 286 mm for the study period of 1985 to 2007). We did not find decrease trend in spring and summer precipitation during the study period (Figure 19). Summer droughts



are common; most of the precipitation falls in spring and fall, coinciding with the two major growing seasons (Walter and Box 1983). The topography is flat with a mean elevation of -15 m below sea level. Soils reflect past transgressions and regressions of the Caspian Sea, and are characterized by a gradient from sandy aeolian deposits and sandy loams in the southeast of the study area to clay loam in the northwest (Kroonenberg et al. 1997).

Vegetation is represented by the typical semi-desert/grassland associations of the northern Precaspian Plain. The main vegetation associations are shortgrass steppe (*Stipa* spp., *Festuca* spp., *Argopyron* spp., and other graminoids) and sage scrub (*Artemisia* spp., *Kochia prostrata*) (Golub 1994). Annual grasses, such as *Bromus tectorum* and *Poa bulbosa* are part of practically all vegetation communities but with varying cover. The presence of *Bromus tectorum* was already recorded in accounts of early explorers of the region (Kostenkov 1868; Krasnov 1886) and is presumable native to our study area. In the current landscape, *Bromus tectorum* is particularly common on recent burns, a pattern that has also been observed in the Western United States, where it is an invasive (Brooks et al. 2004). Shortgrass steppe is characterized by a short growing season in April and May and rapid senescence in the dry summer. The grasses are well adapted to fire due to the dense bunches which protect meristems and generates abundant fuels. Sagebrush (*Artemisia* spp.) dominated shrublands have less biomass than the grasslands, but a longer growing season, and sometimes exhibit a second vegetation peak in the fall and early winter (Kurina and Belousova 1989). *Artemisia* spp. buds are situated above ground and can be killed or damaged by fires. The lack of fire tolerance by *Artemisia* spp. can lead to its gradual substitution by *Stipa* spp. and other graminoids (Neronov 1998). The primary human land use is grazing by domestic livestock, mainly sheep, and to a lesser extent cows and goats.

## Study period

We studied changes in landcover from 1985 to 1998 and from 1998 to 2007 using Landsat satellite images. Our choice of study period was determined by the goal to include vegetation conditions before the collapse of the USSR, the desire to integrate our findings with information from previous research, and last but not least, the availability of remote sensing data for particular phenological periods.

## Data

### *Remote sensing data and image selection*

We used 30-m Landsat 5/7 TM/ETM+ data that correspond to World Reference System (WRS-2) path/row 170/28 for the landcover change classification. Terrain corrected (level 1T) images were acquired from USGS (Figure 25). No additional geometric or radiometric correction was applied. GIMMS AVHRR NDVI and MODIS MOD13A1 data were analyzed to obtain phenology information which aided Landsat image selection. To generate typical vegetation phenology curves, we used Normalized Difference Vegetation Index (NDVI) data from the MOD13A1 dataset, 250-m resolution, 16-day composites for 2007.

Landsat image selection was driven by the observation that the main vegetation communities had different annual phenology curves (Figure 21). Field data and literature indicated that there are at least two periods of spring greenness and these periods are specific for different vegetation types (Matyashenko 1985). Areas that showed the earliest green-up in spring were typically dominated by annual grasses, such as *Bromus tectorum*, and forbs. Areas that greened up later in the spring were typically dominated by perennial grasses, such as *Stipa* spp. Areas dominated by shrubs had a less pronounced spring green-up. For every study year, we derived a curve of the mean NDVI for the entire study area from either GIMMS (1985, 1998) or

MOD13 (2007) data and every potential Landsat image candidate for classification was checked to see if it was right on, or close to, the greenness peak.

Ultimately we selected 2 images (early and late spring) for each study year. Optimal images were available for 1985 and 1998, but no cloud-free Landsat images were available for both phenological periods of 2007. In 2007 the only early spring image available was a suboptimal image because it was recorded in very early spring (March, 10) with portion of the data missing. Landsat 7 experienced difficulty since May, 2003 with its Scan Line Corrector (SLC), resulting in images with gaps that cover 22% of the entire scene (so called SLC-off data) (Maxwell et al. 2007). However, we included this image since there was no alternative.

### *Fire data*

Broad-scale burning in the region can be traced back at least as far as the 18<sup>th</sup> century (Pallas and Blagdon 1802). Though fire was rare in Soviet times, our previous remote sensing analysis showed a sharp increase in burning starting in 1997-1998 which continued until 2007 covering large portion of the study area (Dubinin et al. 2010b).

We used augmented burned areas database from Dubinin et al. 2010b to cover a larger area. Burned areas were delineated based on MODIS MOD02QKM Level 1B calibrated radiance data by visual image interpretation, which is at least as precise as automatic methods (Bowman et al. 2003). Delineating burned areas was straightforward due to distinct fire scar boundaries, often confined by linear features such as roads. The easy recognition of fires was further facilitated by the lack of other disturbances in the area that could have resulted in comparable patterns. Burned areas remained visible for 2 to 3 years after burning, unless a new fire overrode an older scar (Dubinin et al. 2010b).

### *Field data*

Field data were collected during two field seasons in August of 2007, and in May and June of 2008. We collected information on vegetation and disturbances for 160 90-m transects. On each transect, we registered plant species, and litter or bare ground at 270 points that were 0.3 m spaced. We assigned each point to a broad vegetation group based on the phenology of the dominant plant species. The first group represented shrub-dominated communities, which have typically a longer, but less pronounced vegetation peak, and included *Artemisia* species such as *Artemisia lercheana*, *A. austriaca*, *A. pausiflora*, *A. tcherneviana*, as well as *Kochia prostrata*. The second group included perennial grasses of both *Stipa* and *Argopyron* genus, such as: *S. capillata*, *S. sareptana*, *S. lessingiana*, *Agropyron desertorum*, *A. fragile* and others. The third group represented annual grasses, characterized by an early, and typically short, single vegetation peak and included *Bromus tectorum*, *Poa bulbosa*, *Carex arenaria* and others. We defined ‘dominated’ as at least 30% of the live vegetation cover represented by species from this community.

### *Legacy croplands data*

In the 1980s, there were large areas of aeolian dunes which largely corresponded to areas that were either heavily overgrazed or plowed in 1960s and abandoned soon afterwards (Zonn 1995a). About 10% of Republic of Kalmykia was converted to true desert by the early 1980s (Dregne et al. 1991). We digitized the boundaries of permanent sands in 1983 based on natural forage grounds map which had been created in Soviet times using remote sensing data (Trofimov et al. 1983). After abandonment, these formerly plowed were fields first converted to open sands

due to continuing overgrazing, but then started to recover in part due restoration activities initiated by the government and carried out using *Agriophyllum squarrosum*, *Leymus racemosus* as long as variety of annual grasses, and in part, because grazing pressure subsided.

## Analysis

### *Classification*

Previous studies have used multitemporal satellite imagery from specific phenological periods to map changes in vegetation composition (Singh and Glenn 2009), agricultural areas (Lo et al. 1986; Prishchepov et al. 2010), and the phenological heterogeneity of grassland communities (Fisher et al. 2006). Vegetation change can be assessed either via vegetation indices, spectral mixture analysis (Asner and Heidebrecht 2002; Hostert et al. 2003; Kuemmerle et al. 2006), of the classification into different vegetation classes.

We classified the Landsat images into four classes: ‘grass’, ‘shrub’, ‘bare’, and ‘water’. Training data for a supervised classification were collected via a visual interpretation of the images. The ‘grass’ class was identified based on the greenness level, and had digital number values in infrared that were higher than those in red. The ‘bare’ class was defined by high digital number values in all spectral bands, ‘water’ by low digital numbers in all bands, and ‘shrub’ was defined as the remaining non-green areas. We collected at least 200 training points for each class per image.

Each image was classified separately using a Support Vector Machine (SVM). SVM is a machine-learning classification approach that is superior to other classifiers (Foody and Mathur 2004; Huang et al. 2002). SVM is a non-parametric classifier that achieves optimal class separation of classes by fitting hyperplane based on training data. SVM is well suited to map multimodal classes, which parametric-based classifiers (e.g., maximum likelihood classifier)

have difficulty to classify accurately, due to violation of the normality assumption. We used the ImageSVM implementation of SVM (Rabe et al. 2009). We tested a range of parameter combinations ( $g$  from 0.1 to 1000 and  $C$  from 0.1 to 1000) by fitting individual SVM to each parameter pair and comparing models based on three-fold cross-validation thus selecting least-error parameter combination for each image (Rabe et al. 2009). Areas covered with clouds in any Landsat image, and areas of no data due to SLC-off problems were masked out of the resulting classifications of all Landsat images.

The resulting classifications for early and late spring in a given year were combined together to form 9 new vegetation classes. Depending on whether grass-related greenness was mapped in the early or the late spring Landsat image, we defined classes that were dominated by either annuals, perennials, shrubs or permanent sands as well as their mixtures (Table 9). The resulting classes were: ‘annuals-perennials’ – mixed grasslands that include both annuals and perennials in proportions substantial enough to generate green response in early or late spring image, ‘shrub-perennials’ – vegetation communities co-dominated by shrubs and perennial grasses that have a single greenness peak during late spring, ‘perennials’ – vegetation communities characterized by bare ground response during early spring and covered by perennial grasses during late spring, ‘annuals-shrubs’ – communities that have early peak in greenness by lack one afterwards, ‘shrubs’ – areas that do not have neither greenness peaks, nor signs of bare ground, ‘degraded’ – early shrubs and late bare ground and vice versa, ‘annuals’ – early greenness followed by bare ground, and finally, ‘bare’ – bare ground during both periods and ‘water’ – water and any other class during any period.

### *Comparisons*

We overlaid the resulting land cover classifications with the boundaries of bare ground from the Natural forage grounds map of 1983. We then calculated the percentage of correspondence between the bare ground class from the thematic map and the three perennial grasslands classes in 1998 and 2007.

We have compared the resulting land cover classifications with the boundaries of burned areas. We combined annual burned area maps from 1998 to 2007, calculated the total area burned, and mapped burn frequency, which ranged from once to seven times burned from 1998 to 2007. We then calculated: a) the percentage of the entire burned area that occurred in to each land cover class, and b) the percentage of the land cover class that had burned. We also calculated the proportion and area of each burned frequency class in each vegetation class.

## ***Results***

### **Greenness and landcover types**

Field data collected in the spring of 2008 indicated that landcover types dominated by annuals and perennials had higher reflectance in infrared and lower reflectance in red band compared to areas dominated by shrubs, which had less vegetative cover overall (two-tailed T-test,  $N_{\text{annuals}} = 26$ ,  $N_{\text{perennials}} = 98$ ,  $N_{\text{shrubs}} = 51$ , RED: shrubs vs annuals:  $p < 0.01$ , shrubs vs perennials:  $p < 0.01$ ; INFRARED: shrubs vs annual:  $p = 0.03$ , shrubs vs perennials:  $p < 0.01$ , Figure 22). Field samples of perennials and annuals were not significantly different in terms of their brightness (RED:  $p = 0.89$ ; INFRARED:  $p = 0.75$ , Figure 22).

### **Landcover changes**

According to our analysis, the study area experienced a strong increase of grasslands, at the expense of permanent sands and shrubs (Figure 26). Areas covered by perennial grasses more

than doubled from 1985 to 2007 and in the total area of bare ground in 2007 was only 20% of the area they covered in 1985. The increase in grasslands was moderate from 1985 to 1998, the majority of the grassland increase occurred from 1998 to 2007.

Comparisons of the classifications for 1985, 1998, and 2007 showed numerous vegetation changes. For example, only 32% of the area of 'annuals-perennials' class in 1985 stayed in that class by 1998, while 27% transitioned to the 'annuals-shrubs' class. Partly because they were so widespread, the shrub-dominated classes were the most stable (>40%). Transitions from 1985 to 1998 were dominated by transitions from 'shrub' class to 'annuals-shrub'. However, from 1998 to 2007, substantial parts of shrub-dominated classes (36% of 'shrubs-perennials' and 21% of 'shrubs') transitioned to grasses-dominated classes, especially perennials.

Vegetation changes exhibited strong spatial patterns. The most dramatic increase in grasslands coincided with the core area of the Black Lands (Chernye Zemli) where several protected areas have been established since 1991. The pastures in the northern part of the study area, still actively grazed, did not show as much of an increase in perennials, but changes from permanent sands to shrubs and annuals were still common.

### **Association with permanent sands and burning**

Areas that changed from shrubs to grasslands were tightly associated with agricultural fields that were plowed in 1950s and '60s and later abandoned. About half of the areas (42%) mapped as permanent sand in 1985 were converted to one of the three grassland classes (annuals-perennials, shrubs-perennials, perennials) by 1998, and 74% by 2007 (Figure 24). The remaining part of permanent sands was covered by shrublands. However, new grasslands occurred not only on former permanent sands. In addition to perennial grasslands on former permanent sands (450



km<sup>2</sup> in 1985), there were 2491 km<sup>2</sup> areas covered by perennial grasslands in place of shrubs in 1998 and 3528 km<sup>2</sup> in 2007 (Table 12).

Areas of change to perennial grasslands were also associated with burning. However, our analysis could not distinguish if areas that burned were more likely to transition towards perennial grasslands, or if areas that had transitioned to grasslands were more likely to burn. From 1998 until 2007, 38% of the study area burned at least once. Of the burned area 49% was represented by the three perennials classes (which covered 32% of the study area) (Table 10). At the same time 58% of total area of the three perennials classes in 2007 burned at least once between 1998 and 2007, compared to only 38% burned areas for the study area as a whole. Even more striking, the pure perennial class (grass in late spring, and bare ground in early spring), although occupying only a relatively small part of total area burned (7%), burned almost entirely (93%). The three classes dominated by perennial grasses were particularly common in areas where fire frequency was high (Table 11). 65% and more of the areas that burned 3 or more times had perennial grasslands (such areas represented 41% of total burned area and 16% of total study area). The proportion of the burned area represented by perennial grasslands classes increased with increased frequency of burning, with the biggest change between areas that burned twice (45% represented by perennial grasslands) versus 3 times (65% represented by perennial grasslands). On the opposite, classes dominated by shrubs (annuals-shrubs and shrubs classes combined) represented 60% of areas that burned only once, but only 25% of areas that burned twice and less than 10% of the areas that burned 3 times (Table 11).

## ***Discussion***

How human-induced disturbance, and especially grazing affects arid environments is hotly debated. Some studies reported sudden changes, and “catastrophic vegetation shifts”

(Scheffer et al. 2001; van de Koppel et al. 1997) while others argued that recent grazing history has generally little effect on vegetation in arid environments (Milchunas and Lauenroth 1993) compared to other factors such as droughts, fires, insect outbreaks, and floods (Warren et al. 1996; Westoby et al. 1989). As a result, little is known about the ability of semi-desert vegetation to recover once disturbance subsides. Our results suggest that a strong decrease in grazing pressure can indeed lead to substantial changes in dominant species, including a shift from shrublands to grasslands, within a relatively short timeframe (9-10 years) due to changes in both, the direct impact of grazing and alteration of fire regimes.

The breakdown of the Soviet Union overturned dramatic forecasts for the Republic of Kalmykia which was predicted to become completely desertified in 20 years if trends of the 1980s would continue (Vinogradov 1995; Zonn 1995a). In stark contrast, the 90% decrease in livestock has led to an almost complete re-vegetation of open sands by 1998 and a four-fold increase in the area covered by perennial grasslands at the expense of areas previously dominated by shrubs by 2007. This increase in perennial grasses at the expense of shrubs is supported by results from field plots conducted both locally (Neronov 1998) and in the other parts of the world (Grigulis et al. 2005; Hejmanova et al.; Zhang et al. 2005).

Besides decrease in grazing, the increase of grasses could have been potentially caused by several other processes, such as: 1) Increase in atmospheric CO<sub>2</sub>, 2) Climate change, 3) Change in burning regime. Increase in atmospheric CO<sub>2</sub> is a global phenomenon, shown to be beneficial for at least several species of annual grasses (Smith et al. 2000). However, experimental studies highlight that the response to elevated CO<sub>2</sub> level is different for different species and overall response may be negligible (Grunzweig and Körner 2001). Besides grass species like *Bromus tectorum* (Smith et al. 2000), shrubs may also benefit from increase in CO<sub>2</sub>, and global shrub

encroachment is argued to be caused by CO<sub>2</sub> increase as well (Archer et al. 1995). However, Smith et al. observed statistically significant differences between native grasses and *Larrea tridentate* and invasive *Bromus tectorum* at 500 ppm level, which is yet unreachd. Since 1985, the CO<sub>2</sub> concentration in the atmosphere grew at about 1.5 – 2.0 ppm/year rate globally (345 ppm in 1985, and 380 in 2007) (Tans 2010). Given these relatively minor CO<sub>2</sub> concentration increases, and given that they may benefit both grasses and shrubs we suggest that they cannot explain the observed vegetation changes in our study area.

Climate change is another potential driver of vegetation change, and increased precipitation during local and global climatic events, such as El-Nino, can lead to amplified pulses of productivity, especially in annual grasses (Bradley and Mustard 2005; Holmgren et al. 2006). Given the limited availability of cloud-free satellite images, we were not able to fully control for precipitation when selecting study years. However, meteorological data from a station close to the center of the study region (Komsomolsk) did not show substantial differences among the years when our satellite images were taken (92, 125, 95 mm of rainfall total, for March through June of 1985, 1998, and 2007 respectively, Figure 19), and there was no systematic trend in precipitation over the last 25 years. Over next 50 year, a doubling of CO<sub>2</sub> may lead to increases in temperatures of 1-3 C°, and that would lead to higher potential evapotranspiration, which, assuming that rainfall will not change, will increase aridity. However, photosynthesis boost from increasing CO<sub>2</sub> would also increase water-use efficiency via the reduction of stomatal conductance making it difficult predicting net balance. In any event though, the observed recent vegetation changes cannot be attributed to climate change. Thus, changes in CO<sub>2</sub> concentration, temperature and precipitation may be by far outweighed by direct human impacts (Amthor

1995; LeHouérou 1996). Indeed, compared with slowly changing background factors, changes in burning and land use are more pronounced and rapid (Dubinin et al. 2010a).

The increase in grasslands may indicate a positive feedback loop given the substantial association between patterns of fire and shifts to grasslands in our study area. Grasslands provide more fine-fuel easily flammable material than shrub-dominated communities (D'Antonio and Vitousek 1992). Furthermore, grasses can recover easier after fire, which thus creates a competitive advantage over shrubs (Grigulis et al. 2005).

It is not clear if the observed fire patterns, and hence the expansion of grasslands, are 'natural'. Historical literature provides mixed evidence on extent and frequency of fires in the North-West Caspian lowland in the 19<sup>th</sup> and 20<sup>th</sup> centuries. Some sources mention widespread grassland fires (Pallas and Blagdon 1802; Zwick and Schill 1831). However, reports from research expeditions, specifically aimed to study of natural resources and management do not mention any grasslands fires (Kostenkov 1868; Krasnov 1886), but this does not necessarily reflect their actual absence. Based on that we speculate that the release of livestock pressure in 1990s and restoration of fuel loads might have reverted vegetation communities to pre-Soviet conditions overall, potentially with grasses being more widespread. However, given the rebounding trend of livestock since 1999, we expect a potential decline in fires and a return of shrubs in the near future.

While fire is certainly the most important driver of vegetation change, it is not the only cause of the increase in grasslands. We found that a substantial proportion of the areas that shifted to grass-dominated communities had not burned. On the other hand, not all areas that did burn, converted to grasslands, suggesting interaction with soil type, fire return interval and exact timing of burning and land use legacies. In general, bare grounds undergoing first stages of

succession were more likely to burn, as they are first populated by annuals. Additionally restoration efforts also led to establishment of species with high fuel load and contiguity.

Our satellite classifications did not provide species-level change information, and mapped only broad phenological groups. However, it is of interest to try to compare the role of annual grasses, especially *Bromus tectorum* L., in the study area with its role in other areas of the world, where it represents highly invasive species and can lead to change in fire dynamics. Cheatgrass is a highly competitive pioneer species in both its native and foreign range (Mack 1981; Neronov 2006). While in invasive mode, cheatgrass can dominate vegetation communities forming monocultures in shrub or bunchgrass ecosystem (Mack 1981), but in its native range it is usually co-exist with perennial grasses and shrubs, such as of *Stipa* and *Artemisia*. Our results indeed showed common presence of an early green-up, which we interpreted as an annual grass signal in ‘annuals-perennials’, ‘annuals-shrubs’ classes, and we found very few areas that were pure ‘annuals’. Similarly, our field data indicated that cheatgrass and other annuals were present in low abundances in most of the vegetation communities in our study area (results not shown).

## **Conclusions**

The research that we present here provides evidence for substantial vegetation change in Kalmykia. These vegetation changes are a complex phenomenon, and were probably caused by a multitude of factors. The sharp increase in annual burned area in the semi-deserts of Southern Russia is most likely the main direct driving force of vegetation change in the region. However, burning in turn is affected by human activities such as livestock grazing (Dubinin et al. 2010a). Land use thus can impact the structure and composition of vegetation communities and outweighed in our study area the effects of increasing CO<sub>2</sub> concentration and climatic change.

## ***Acknowledgements***

We gratefully acknowledge support for this research by the NASA Land-Cover and Land-Use Change Program (NNG05GA74G), by a NASA Earth System Science Dissertation Fellowship for M. Dubinin (07-Earth07F-0053), by the U.S. Fish and Wildlife Service, and by a grant from the National Geographic Society (8360-07). We are also grateful to field assistants A. Zakharov, S. Goryunova, E. Degtyarev, I. Glushkov.

## ***Bibliography***

- Amthor, J.S. (1995). Terrestrial Higher-Plant Response to Increasing Atmospheric [CO<sub>2</sub>] in Relation to the Global Carbon-Cycle. *Global Change Biology*, 1, 243-274
- Archer, S. (1994). Woody plant encroachment into Southwestern grasslands and savannas; rates, patterns and proximate causes. In M. Vavra, W. Laycock & R. Pieper (Eds.) *Ecological implications of livestock herbivory in the west* (pp. 13-68). Denver, CO Soc. Range Manage.
- Archer, S., Schimel, D.S., & Holland, E.A. (1995). Mechanisms of shrubland expansion - land-use, climate or CO<sub>2</sub>. *Climatic Change*, 29, 91-99
- Asner, G.P., & Heidebrecht, K.B. (2002). Spectral unmixing of vegetation, soil and dry carbon cover in arid regions: comparing multispectral and hyperspectral observations. *International Journal of Remote Sensing*, 23, 3939-3958
- Belsky, J.A. (1992). Effects of Grazing, Competition, Disturbance and Fire on Species Composition and Diversity in Grassland Communities. *Journal of Vegetation Science*, 3, 187-200
- Bond, W.J., Woodward, F.I., & Midgley, G.F. (2005). The global distribution of ecosystems in a world without fire. *New Phytologist*, 165, 525-537
- Bowman, D., Zhang, Y., Walsh, A., & Williams, R.J. (2003). Experimental comparison of four remote sensing techniques to map tropical savanna fire-scars using Landsat-TM imagery. *International Journal of Wildland Fire*, 12, 341-348
- Bradley, B.A., & Mustard, J.F. (2005). Identifying land cover variability distinct from land cover change: Cheatgrass in the Great Basin. *Remote Sensing of Environment*, 94, 204-213

- Brooks, M.L., D'Antonio, C.M., Richardson, D.M., Grace, J.B., Keeley, J.E., DiTomaso, J.M., Hobbs, R.J., Pellant, M., & Pyke, D. (2004). Effects of invasive alien plants on fire regimes. *Bioscience*, 54, 677-688
- Callaway, R.M., & Davis, F.W. (1993). Vegetation Dynamics, Fire, and the Physical-Environment in Coastal Central California. *Ecology*, 74, 1567-1578
- CIESIN, & CIAT (2005) Gridded Population of the World Version 3 (GPWv3): Population Density Grids. Palisades, NY; Available at <http://sedac.ciesin.columbia.edu/gpw>. Last accessed: 06.28.2009: Socioeconomic Data and Applications Center (SEDAC), Columbia University
- CIESIN, FAO, & CIAT (2005) Gridded Population of the World: Future Estimates, 2015 (GPW2015): Population Density Grids. Palisades, NY; Available at <http://sedac.ciesin.columbia.edu/gpw>. Last accessed: 06.28.2009: Socioeconomic Data and Applications Center (SEDAC), Columbia University
- D'Antonio, C.M., & Vitousek, P.M. (1992). Biological Invasions by Exotic Grasses, the Grass Fire Cycle, and Global Change. *Annual Review of Ecology and Systematics*, 23, 63-87
- Dregne, H., Kassas, M., & Rozanov, B. (1991). A new assessment of the world status of desertification. *United Nations Environment Programme Desertification Control Bulletin*, 20, 6-18
- Dubinin, M., Lushchekina, A., & Radeloff, V.C. (2010a). Relationships between climate, livestock, and fire in the arid grasslands of southern Russia (in review)
- Dubinin, M., Potapov, P., Lushchekina, A., & Radeloff, V.C. (2010b). Reconstructing long time series of burned areas in arid grasslands of southern Russia by satellite remote sensing. *Remote Sensing of Environment*, 114, 1638-1648
- FAO (2010) FAOSTAT, <http://faostat.fao.org>



- Fisher, J.I., Mustard, J.F., & Vadeboncoeur, M.A. (2006). Green leaf phenology at Landsat resolution: Scaling from the field to the satellite. *Remote Sensing of Environment*, 100, 265-279
- Foody, G.M., & Mathur, A. (2004). A relative evaluation of multiclass image classification by support vector machines. *IEEE Transactions on Geoscience and Remote Sensing*, 42, 1335-1343
- Foster, D.R., Knight, D.H., & Franklin, J.F. (1998). Landscape patterns and legacies resulting from large, infrequent forest disturbances. *Ecosystems*, 1, 497-510
- Golub, V.B. (1994). The desert vegetation communities of the Lower Volga Valley. *Feddes Repertorium*, 105, 499-515
- Grigulis, K., Lavorel, S., Davies, I.D., Dossantos, A., Lloret, F., & Vila, M. (2005). Landscape-scale positive feedbacks between fire and expansion of the large tussock grass, *Ampelodesmos mauritanica* in Catalan shrublands. *Global Change Biology*, 11, 1042-1053
- Grunzweig, J.M., & Korner, C. (2001). Growth, water and nitrogen relations in grassland model ecosystems of the semi-arid Negev of Israel exposed to elevated CO<sub>2</sub>. *Oecologia*, 128, 251-262
- Hejermanova, P., Hejerman, M., Camara, A.A., & Antoninova, M. Exclusion of livestock grazing and wood collection in dryland savannah: an effect on long-term vegetation succession. *African Journal of Ecology*, 48, 408-417
- HilleRisLambers, J., Yelenik, S.G., Colman, B.P., & Levine, J.M. (2010). California annual grass invaders: the drivers or passengers of change? *Journal of Ecology*, 98, 1147-1156
- Hoelzel, N., Haub, C., Ingelfinger, M.P., Otte, A., & Pilipenko, V.N. (2002). The return of the steppe: Large-scale restoration of degraded land in southern Russia during the post-Soviet era. *Journal for Nature Conservation (Jena)*, 10, 75-85
- Holmgren, M., Stapp, P., Dickman, C.R., Gracia, C., Graham, S., Gutierrez, J.R., Hice, C., Jaksic, F., Kelt, D.A., Letnic, M., Lima, M., Lopez, B.C., Meserve, P.L., Milstead, W.B., Polis, G.A.,

- Previtali, M.A., Michael, R., Sabate, S., & Squeo, F.A. (2006). Extreme climatic events shape arid and semiarid ecosystems. *Frontiers in Ecology and the Environment*, 4, 87-95
- Hostert, P., Roeder, A., Hill, J., Udelhoven, T., & Tsiourlis, G. (2003). Retrospective studies of grazing-induced land degradation: a case study in central Crete, Greece. *International Journal of Remote Sensing*, 24, 4019-4034
- Huang, C., Davis, L.S., & Townshend, J.R.G. (2002). An assessment of support vector machines for land cover classification. *International Journal of Remote Sensing*, 23, 725-749
- Kostenkov, K. (1868). Statistics and management description of Kalmyk steppe. *Kalmyk steppe of Astrakhan region, study by Kumo-Manych expedition*. Saint-Petersburg Ministry of State Property
- Krasnov, F. (1886). *Geobotanic research in Kalmykian steppes*. Saint-Petersburg (in Russian)
- Kroonenberg, S.B., Rusakov, G.V., & Svitoch, A.A. (1997). The wandering of the Volga delta: A response to rapid Caspian sea-level change. *Sedimentary Geology*, 107, 189-209
- Kuemmerle, T., der, A., & Hill, J. (2006). Separating grassland and shrub vegetation by multivariate pixeladaptive spectral mixture analysis. *International Journal of Remote Sensing*, 27, 3251-3271
- Kurinova, Z.S., & Belousova, Z.N. (1989). *Effective use of forage resources*. Elista: Kalmyk book publishing
- LeHouerou, H.N. (1996). Climate change, drought and desertification. *Journal of Arid Environments*, 34, 133-185
- Lo, T.H.C., Scarpace, F.L., & Lillesand, T.M. (1986). Use of Multitemporal Spectral Profiles in Agricultural Land-Cover Classification. *Photogrammetric Engineering and Remote Sensing*, 52, 535-544

- Mack, R.N. (1981). Invasion of *Bromus tectorum* L. into Western North America - an ecological chronicle. *Agro-Ecosystems*, 7, 145-165
- Matyashenko, G.V. (1985). *Vegetation dynamics of chalk deposits of pre Ural plateau [in Russian]*. Novosibirsk: Nauka
- Maxwell, S.K., Schmidt, G.L., & Storey, J.C. (2007). A multi-scale segmentation approach to filling gaps in Landsat ETM+ SLC-off images. *International Journal of Remote Sensing*, 28, 5339-5356
- Milchunas, D.G., & Lauenroth, W.K. (1993). Quantitative Effects of Grazing on Vegetation and Soils over a Global Range of Environments. *Ecological Monographs*, 63, 327-366
- Neronov, V.V. (1998). Antropogenic grasslands encroachment of desert pastures of North-Western Precaspian plains. *Successes of modern biology*, 118, 597-612 (in Russian)
- Neronov, V.V. (2006). Dynamics of biological diversity of the degraded arid ecosystems during their natural restoration (after the example of Chernye Zemli, Kalmykia) (in Russian). *Izvestiya Akademii Nauk Seriya Geographicheskaya*, 111-119
- Olden, J.D., Poff, N.L., Douglas, M.R., Douglas, M.E., & Fausch, K.D. (2004). Ecological and evolutionary consequences of biotic homogenization. *Trends in Ecology & Evolution*, 19, 18-24
- Pallas, P.S., & Blagdon, F.W. (1802). *Travels through the southern provinces of the Russian Empire, in the years 1793 and 1794*. London: Printed by A. Strahan ... for T.N. Longman and O. Rees ...
- Prishchepov, A.V., Radeloff, V.C., Dubinin, M., & Alcantara, C. (2010). The effect of image acquisition dates on detection of agricultural land abandonment in Eastern Europe (in review). *Remote Sensing of Environment*
- Rabe, A., van der Linden, S., & Hostert, P. (2009). ImageSVM 2.0

- Roques, K.G., O'Connor, T.G., & Watkinson, A.R. (2001). Dynamics of shrub encroachment in an African savanna: relative influences of fire, herbivory, rainfall and density dependence. *Journal of Applied Ecology*, 38, 268-280
- ROSSTAT (2006). All-russian agricultural survey. Livestock population. [in Russian]. Available at: <http://www.gks.ru/news/perepis2006/totals-osn.htm>, 5, 38-39
- Scheffer, M., Carpenter, S., Foley, J.A., Folke, C., & Walker, B. (2001). Catastrophic shifts in ecosystems. *Nature*, 413, 591-596
- Schlesinger, W.H., Reynolds, J.F., Cunningham, G.L., Huenneke, L.F., Jarrell, W.M., Virginia, R.A., & Whitford, W.G. (1990). Biological Feedbacks in Global Desertification. *Science*, 247, 1043-1048
- Shilova, S.A., Neronov, V.V., Kasatkin, M.V., Savinetskaya, L.E., & Tchabovsky, A.V. (2007). Fires at the current stage of evolution of semi-desert in southern Russia: the influence on vegetation and rodent populations. *Uspekhi Sovremennoi Biologii*, 127, 372-386
- Singh, N., & Glenn, N.F. (2009). Multitemporal spectral analysis for cheatgrass (*Bromus tectorum*) classification. *International Journal of Remote Sensing*, 30, 3441-3462
- Smith, S.D., Huxman, T.E., Zitzer, S.F., Charlet, T.N., Housman, D.C., Coleman, J.S., Fenstermaker, L.K., Seemann, J.R., & Nowak, R.S. (2000). Elevated CO<sub>2</sub> increases productivity and invasive species success in an arid ecosystem. *Nature*, 408, 79-82
- Tans, P. (2010) Globally averaged marine surface monthly mean data. NOAA/ESRL ([www.esrl.noaa.gov/gmd/ccgg/trends](http://www.esrl.noaa.gov/gmd/ccgg/trends)).
- Trofimov, I.A., Vyshivkin, D.D., & Savchenko, I.V. (1983) Natural forage grounds. 1:500000 *Natural ecosystems of Kalmyk ASSR*. V.P. Williams Forage Institute.

- Van Auken, O.W. (2000). Shrub invasions of North American semiarid grasslands. *Annual Review of Ecology and Systematics*, 31, 197-215
- Van Auken, O.W. (2009). Causes and consequences of woody plant encroachment into western North American grasslands. *Journal of Environmental Management*, 90, 2931-2942
- van de Koppel, J., Rietkerk, M., & Weissing, F.J. (1997). Catastrophic vegetation shifts and soil degradation in terrestrial grazing systems. *Trends in Ecology & Evolution*, 12, 352-356
- van Langevelde, F., van de Vijver, C., Kumar, L., van de Koppel, J., de Ridder, N., van Andel, J., Skidmore, A.K., Hearne, J.W., Stroosnijder, L., Bond, W.J., Prins, H.H.T., & Rietkerk, M. (2003). Effects of fire and herbivory on the stability of savanna ecosystems. *Ecology*, 84, 337-350
- Vinogradov, B.V. (1995). Forecasting dynamics of open sands of Black Lands, Kalmykia using repeated aerial photography. In N. Zonn I.S., V.M. (Ed.) *Biota and Environment of Kalmykia* (pp. 259-268). Moscow, Elista (in Russian)
- Walter, H., & Box, E. (1983). Overview of Eurasian continental deserts and semideserts. *Ecosystems of the World* (pp. 3-269). Amsterdam Elsevier
- Warren, A., Sud, Y.C., & Rozanov, B. (1996). The future of deserts. *Journal of Arid Environments*, 32, 75-89
- Westoby, M., Walker, B., & Noymeir, I. (1989). Range Management on the Basis of a Model Which Does Not Seek to Establish Equilibrium. *Journal of Arid Environments*, 17, 235-239
- Zarin, D.J., Davidson, E.A., Brondizio, E., Vieira, I.C.G., Sa, T., Feldpausch, T., Schuur, E.A., Mesquita, R., Moran, E., Delamonica, P., Ducey, M.J., Hurtt, G.C., Salimon, C., & Denich, M. (2005). Legacy of fire slows carbon accumulation in Amazonian forest regrowth. *Frontiers in Ecology and the Environment*, 3, 365-369

- Zhang, J.Y., Wang, Y., Zhao, X., & Zhang, T. (2005). Grassland recovery by protection from grazing in a semi-arid sandy region of northern China. *New Zealand Journal of Agricultural Research*, 48, 277-284
- Zonn, I.S. (1995a). Desertification in Russia: Problems and solutions (an example in the Republic of Kalmykia-Khalmg Tangch). *Environmental Monitoring and Assessment*, 37, 347-363
- Zonn, S.V. (1995b). Republic of Kalmykia - Khalmg Tangch - European region of environmental tension. *Biota and Environment of Kalmykia* (pp. 6-18). Moscow, Elista (in Russian)
- Zwick, H.A., & Schill, J.G. (1831). *Calmuc Tartary or journey from Sarepta to several Calmuc Hordes of the Astrakhan Government*. London: Holdsworth and Ball

## Tables

Table 9. The definition of nine vegetation associations and other land cover classes for each year based on the classification of four land cover classes for corresponding early and a late spring classifications.

		Early spring classification			
		grass	shrub	bare ground	water
Late spring classification	grass	annuals-perennials	shrubs-perennials	perennials	water
	shrub	annuals-shrubs	shrubs	degraded	water
	bare				
	ground	annuals	degraded	permanent sand	water
	water	water	water	water	water

Table 10. The association of different land cover classes in 2007 with the cumulative burned areas (1998 – 2007). Subtotal for perennial grasslands is the sum of the three classes that had perennials (annuals-perennials, shrubs-perennials, perennials), the subtotal for shrubs is the sum of two shrub classes (annuals-shrubs, shrubs).

		% of the	Burned		% of the total area in this vegetation class that burned
	Total area (km <sup>2</sup> )	total area	area (km <sup>2</sup> )	% of the total burned area	
Annuals-perennials	4,320,863	21%	2,433,805	32%	56%
Shrubs-perennials	1,567,060	8%	754,644	10%	48%
Perennials	618,161	3%	572,526	7%	93%
Annuals-shrubs	4,979,386	24%	978,850	13%	20%
Shrubs	6,424,930	31%	1,989,557	26%	31%
Degraded	422,474	2%	216,439	3%	51%
Annuals	11,424	0%	1,244	0%	11%
Permanent sand	278,290	1%	60,991	1%	22%
Subtotal for perennials	6,506,084	32%	3760975	49%	58%
Subtotal for shrubs	11,404,316	55%	2968407	39%	26%



Table 11. The proportion of each land cover class that burned once, twice, etc. up to seven time.

	Number of times an area burned from 1998 to 2007						
	1	2	3	4	5	6	7
Annuals-perennials	19%	28%	40%	44%	53%	44%	49%
Shrubs-perennials	7%	10%	12%	12%	12%	17%	7%
Perennials	3%	7%	13%	12%	9%	9%	11%
Annuals-shrubs	20%	13%	8%	4%	3%	3%	2%
Shrubs	40%	30%	13%	9%	6%	9%	7%
Degraded	3%	3%	2%	3%	3%	2%	6%
Annuals	0%	0%	0%	0%	0%	0%	0%
Permanent sand	1%	1%	0%	0%	0%	0%	0%
Subtotal for							
perennials	30%	45%	65%	69%	74%	70%	67%
Subtotal for shrubs	63%	46%	22%	17%	13%	14%	15%
Total burned area,							
(km <sup>2</sup> )	2628	1538	1221	931	482	189	29





## Figures

Figure 17. Average annual decline in total sheep from 1992 to 2008 for the ten countries with the strongest absolute declines. Number in brackets indicates time span in years between the lowest and the highest sheep number during this period (FAO 2010).




Figure 18. Study area location (hatched area with black outline) in Southern Russia.



Figure 19. Trends in spring NDVI and precipitation (sum of March-June period rainfall, mm) in the study area from 1982 to 2007 period. While there were strong increases in NDVI, there was no systematic trend in precipitation.




Figure 20. Examples of different vegetation types, from top left clockwise: *Stipa capillata* dominated community, *Artemisia lercheana* dominated community, community dominated by annual *Bromus tectorum* and perennial *Stipa sareptana*, permanent bare sands. Images taken by M. Dubinin in late spring.



Figure 21. Phenological differences for the three pure vegetation types based on field data and MOD13A1 NDVI in 2007, error bars indicate 1 standard deviation ( $N_{\text{perennials}} = 6$ ,  $N_{\text{shrubs}} = 6$ ,  $N_{\text{annuals}} = 6$ ).






Figure 22. Differences in brightness (0-255, Landsat 5/TM May 5, 2007) for different vegetation communities,  $N_{\text{shrubs}} = 51$ ,  $N_{\text{annuals}} = 26$ ,  $N_{\text{perennials}} = 98$ .




Figure 23. Total area in different land covers classes in 1985, 1998, and 2007.




Figure 24. Association between former permanent sands (red polygons) and landcover (green – grasslands, grey – shrublands, yellow – permanent sands, blue - water) in late spring of 1998 (left, top) and late spring of 2007 (left, bottom). Also shown fragments of Landsat TM images, RGB combination: bands 5, 4, and 3 (right)





Figure 25. Annual NDVI trends for the years of selected satellite images in a) 1985, Landsat images recorded on May 8<sup>th</sup> and June 6<sup>th</sup> and 1998 April 26<sup>th</sup> and June 13<sup>th</sup>, and b) 2007 (March 10<sup>th</sup> and May 5<sup>th</sup>).

a)



b)







Figure 26. Land cover classifications for 1985 (top left), 1998 (top right), and 2007 (bottom right). White areas were masked out because of cloud contamination in at least one of the images. The apparent striping is caused by Landsat problems with the scan line correction procedures which caused stripes of missing data.



## Chapter 4: Saiga habitat selection and habitat distribution in southern Russia

### ***Abstract***

Understanding the habitat preferences of endangered species is important for their long-term conservation. Saiga antelope (*Saiga tatarica tatarica* L., 1766) is the last free-roaming antelope in Europe, and is highly endangered. Saiga antelope numbers plummeted, first in the early 1980s, and again in the mid-1990s following the collapse of the Soviet Union. Though the decline of Saiga antelope is primarily attributed to poaching for its horns, substantial changes in vegetation and increasing fire frequency over the last 20 years may also have affected Saiga antelope habitat selection. Unfortunately, not much is known about Saiga antelope habitat and its selection in general and its change over time. The goal of this study was to understand seasonal Saiga antelope habitat selection and to map habitat changes between 2003 and 2007 in Southern Russia. We analyzed Saiga antelope presence-only data for 2003-2007, collected in two nature reserves in the study region. We used Maxent (Phillips, 2006) to explore the relative importance of a suite of environmental and human-disturbance variables for Saiga habitat selection, and to predict the distribution of Saiga habitat. We constructed a global model for the entire dataset, and models based on subset of our data representing different years and seasons. We hypothesized that vegetation composition is a key factor explaining Saiga presences, while fire-affected areas will be of less interest for Saiga antelope at least in particular seasons. Second, we hypothesized that habitat selection differs significantly among seasons, but is relatively stable among years.

To our surprise we found that vegetation patterns did not explain Saiga habitat selection well, but that Saiga antelopes selected areas that burned more often. Distance to water sources and proximity to farms were also important factors influencing Saiga antelope habitat selection. Variable importance did not vary substantially between years and seasons. Overall, suitable habitat occupied about 20% of the 2,500 km<sup>2</sup> study area, and the amount of suitable habitat did not change substantially during the time period studied. Our results indicate that suitable Saiga habitat was characterized by variables describing environmental change. The importance of burned areas indicates that Saiga antelope appear to select open areas, where poachers and predators can be more seen easily. Though vegetation variables were less important than burned areas and human disturbance variables in our models, the burning frequency variable may be a proxy for productivity of herbs and grasses, and thus forage availability. To ensure the conservation of this species the new understanding of Saiga habitat preferences our study provides should be incorporated into management and conservation practices.

## ***Introduction***

One of the essential questions for the conservation of large ungulates is whether their population declines are caused directly by humans, for example via poaching, or indirectly, for example via habitat loss caused by land-use or climate change (Ehrlich and Ehrlich 1981). Habitat loss and fragmentation are arguably the largest threat for populations of large mammals (Simberloff 1984). Poaching is a second major threat for ungulate populations, because these species are often hunted for their meat, medicinal properties, or trophies (Milner-Gulland and Bennett 2003; Morrison et al. 2007). Poaching is especially widespread in the developing world, where poverty often forces people to exploit wildlife populations to subsidize their diets and incomes. Moreover, where collapsing states or warfare result in an eroding infrastructure for nature protection and a low level of control, poaching can become the main cause of species endangerment (Bhatnagar et al. 2009; Dudley et al. 2002). Habitat change and poaching often co-occur (Sodhi et al. 2004; Wilkie et al. 2000), making it difficult to isolate their relative effect. Habitat selection studies in areas where poaching is restricted (i.e., in protected areas) can reveal how environmental change affects habitat availability, and ultimately may facilitate understanding of past population change.

The Saiga antelope (*Saiga tatarica* spp. *tatarica*) is a migratory species whose population once numbered in the millions and whose range stretched from the Sea of Azov in the West to Mongolia in the East. Currently, saiga antelope remain in substantial yet low numbers only in one part of Russia (North Western Caspian region) and in a few disjunct populations in Kazakhstan and Mongolia (Bekenov et al. 1998; Milner-Gulland et al. 2003). Saiga antelope populations precipitously declined throughout their range first in the beginning of 1980s, and



again after the collapse of the Soviet Union (Figure 27) (Milner-Gulland et al. 2001). When the population decline started, its causes were unclear, with some studies highlighting poaching and others suggesting that skewed sex ratios and environmental degradation were the main cause (Chan et al. 1995; Milner-Gulland et al. 1995). The harsh economic crisis of 1998 resulted in a further decline of living conditions in rural Russia. The radical withdrawal of agricultural subsidies and a lack of alternative economic opportunities resulted in local people turning to poaching Saiga antelope for meat and horns (Robinson 2000), which is now seen as the predominant factor driving Saiga population collapse (Karimova 2002; Milner-Gulland et al. 2001). These drivers of population decline were also consistent among populations in Russia and Kazakhstan (Kuhl et al. 2009a).

The same socio-economic transformations and especially the breakdown of many state-farms in the transition period also led to rapid and widespread environmental changes across the range of Saiga antelope in Kalmykia, mainly spurred by a sharp decrease in numbers of livestock. These environmental changes, however, were lagged in time. Among the documented changes were a steady increase in fire frequency and burned areas since 1980s, with substantially more burning after 1997, covering up to 20% of the Saiga antelope's range in Russia (Dubinin et al. 2010c). Associated with changes in fire regimes, substantial vegetation changes occurred, resulting in an increase of areas covered by perennial grasses at the expense of areas covered by shrubs (Dubinin et al. 2010b). These vegetation changes may have resulted in Saiga habitat loss, potentially contributing to the Saiga antelope decline in substantial ways (Abaturov 2007). Overall, potential Saiga habitat loss is of major concern for the conservation of this species, because it is even harder to control than poaching (Karimova 2002)

During the Soviet period, Saiga antelope were an important natural resource and the species was industrially hunted and therefore the subject of numerous studies. However, these studies were rarely spatially-explicit and usually provided only descriptive analysis of habitat use and diet (Bannikov 1967; Sokolov and Zhirnov 1998). When the population decline started, research almost exclusively focused on population parameters, such as sex ratios, survival, and fecundity (Kuhl et al. 2009b; Milner-Gulland et al. 2003). Though environmental conditions were not a research focus a few studies provide insight into Saiga habitat selection, for example showing that Saiga antelopes avoid perennial grasslands (Larionov et al. 2008). Current research was focused on calving site selection and migrations in Kazakhstan and Mongolia (Berger et al. 2008; Singh et al. 2010a, b). Overall though, what determines suitable Saiga habitat and how Saiga habitat has change recently remains almost unknown.

Our overarching goal was to analyze habitat selection of Saiga antelopes antelope in the Chernye Zemli region of the Republic of Kalmykia, to predict habitat suitability, and to assess how habitat preferences and habitat patterns change among years and seasons. Our specific objectives were: 1) to predict habitat suitability from 2003 until 2008; 2) to understand the role of different environmental and human factors in characterizing suitable habitat; and 3) to determine the inter-annual and seasonal variability of habitat availability and the driving factors of habitat selection. We hypothesized that: 1) vegetation is an important factor for habitat selection of Saiga, 2) there are no significant differences in habitat patterns between years, 3) differences between suitable habitats of different seasons are substantial; and 4) fire affected areas are avoided by Saiga antelope.

## **Methods**

### **Study area**

Our study area occupies 2,500 km<sup>2</sup> of the Republic of Kalmykia and Astrakhan Region in southern Russia (Figure 28). The study area includes two protected areas: Stepnoy Nature Preserve and a part of Chernye Zemli Nature Reserve. Whereas the nature preserve is protected on a periodic watch basis, with one to two rangers teams with motorcycles patrolling the territory, the State Nature Reserve has permanent protection staff.

The climate of the study area is arid, with hot, dry summers. Mean daily temperature is +24°C in July with maximum temperatures of up to +44°C. Annual precipitation is 150 to 350 mm, with most of the precipitation in spring and fall (Walter and Box 1983). The topography is rolling with a mean elevation of -15 m below sea level.

Vegetation is typical for semi-deserts of the northern Precaspian Plains. Dominant vegetation associations are shortgrass steppe (*Stipa* spp., *Festuca* spp., *Argopyron* spp., and other graminoids) and dwarf sage shrublands (*Artemisia* spp., *Kochia prostrata*) with ubiquitous annuals such as *Bromus tectorum* and *Poa bulbosa* (Golub 1994). Shortgrass steppe has short growing season in April and May and rapid senescence in the dry summer. The grasses are protected from fire by dense bunches and resprout with the first rain after fire. Dwarf shrublands have less biomass than the grasslands, but a longer growing season, and sometimes exhibit a second vegetation peak in the fall and early winter (Kurinova and Belousova 1989). The primary human land use is grazing by domestic livestock, mainly sheep, and to a lesser extent cows and goats.

## Data

### *Occurrence data*

We studied changes in habitat selection from 2003 to 2007. Our choice of study period was determined by the availability of Saiga antelope presence data. Herd observation data were collected by the staff of the Chernye Zemli Nature Reserve and Stepnoi Nature Preserve under a protocol developed during an INTAS (International Association for the promotion of co-operation with scientists from the New Independent States of the former Soviet Union, EU) project. Saiga antelopes were observed from a vehicle or on foot while on patrol between 0200 hours and 2100 hours using binoculars. Data collection was opportunistic, with some areas being covered better and more often than others, thus introducing some sampling bias (Phillips et al 2009, see below). Different roads were used by rangers depending on road conditions and passability due to weather conditions.

A total of 1,979 direct observations of individuals and herds collected from September 2003 until December 2007 were analyzed. The maximum distance at which saiga were detected was about 600 meters, with a typical range of 250-600 meters (A. Khludnev, personal communication). To reduce positional uncertainty in our presence data, we therefore generalized the points to a 250-m (binary: presence / background) grid for total, yearly and seasonal models. A second reason for generalizing our dataset was to reduce spatial autocorrelation effects. Generalizing decreased the number of locations to about half of the full dataset (i.e., leaving 851 independent observations, Table 13).

### *Predictor variables*

As a proxy for vegetation productivity, we calculated total, annual, and seasonal (across the entire period of study) mean and standard deviation of the Natural Difference Vegetation Index (NDVI) based on 250-m resolution MOD13A1 (23, 16-day composites per year) images. In addition, we used Landsat-based phenological vegetation classifications for 1998 and 2007 representing nine classes: 1 - mixed grasslands with both annuals and perennials, 2 - vegetation communities co-dominated by shrubs and perennial grasses, 3 – pure perennial grasses, 4 – annual grasses and shrubs, 5 – pure shrubs, 6 – degraded areas, 7 - vegetation communities dominated by annuals, 8 - bare ground and 9 – water (Dubinin et al. 2010a).

To capture burned area, we extended the burned area database from Dubinin et al. (2010b) to cover our entire study area. Burned areas were digitized from satellite data from 1998, when active burning started, until 2008 (Dubinin et al. 2010c) using the RESURS/MSU-SK and TERRA/MODIS MOD02QKM Level 1B calibrated radiance images (Bowman et al. 2003). Delineating burned areas was straightforward due to distinct fire scar boundaries, often created by linear features such as roads. Furthermore, there are no other disturbances in the area that could have resulted in comparable patterns. Burned areas remained visible for 2 to 3 years after burning, unless a new fire overrode an older scar (Dubinin et al. 2010c). For each year of the study period we calculated two fire metrics for each 250-m pixel: 1) a binary classification of burned versus non-burned and 2) the number of times a particular area burned prior to the current year (i.e., in 2003 the variable described the number of times a pixel had burned from 1998 until 2002).

We used the Shuttle Radar Topography Mission (SRTM) elevation model (<http://srtm.csi.cgiar.org>) to obtain absolute elevation and a terrain ruggedness index (Riley et al. 1999). These two measures were used as proxies of disturbance, as Saiga antelope are assumed

to select for higher and more flat terrain (A. Khudnev, personal communication). Human disturbance was measured as the distance to active farms. Initially, we also tested a distance-to-roads variable for inclusion in the models, but removed it because it interfered with our sampling bias in the occurrence data. Since most of the Saiga locations were obtained along roads, this variable caused areas near roads to be identified as optimal habitat, which was ecologically not meaningful.

### **Habitat suitability modelling**

We used a maximum entropy modeling approach to analyze Saiga antelope suitable habitat. Maximum entropy modeling is a machine learning approach that uses species' presence data to predict their spatial distributions (Phillips et al., 2006). The true, but unknown, distribution of a species is approximated by deriving a probability distribution using constraints inferred from environmental variables associated with occurrence data points. The maximum entropy principle suggests that the distribution that approximates the 'true' distribution best given all current knowledge is the distribution with maximum entropy (Jaynes, 1957). To avoid over-fitted models, regularization parameters are used (Phillips et al., 2006; Phillips and Dudik, 2008). Maximum entropy models perform well with small sample sizes (Wisz et al., 2008) and frequently outperform classical statistical approaches in their predictive power (Elith et al. 2006).

To fit maximum entropy models, we used the software Maxent (version 3.3a, <http://www.cs.princeton.edu/~schapire/maxent/>). For all model runs we used a maximum number of 2,500 iterations with 5 replicate runs and default regularization parameters (Phillips and Dudik, 2008). We used 75% of the data to train models and 25% to test them. To estimate Maxent distribution, 10,000 random background points were collected. To account for sampling bias in our Saiga locations (because occurrence points were mostly collected along roads), we

sampled background points only within 1-km buffers around roads (Phillips et al. 2009). The distance of 1 km was chosen because it represented the 50% percentile of the distribution of all presence point distances to roads.

Model validation was based on the area under the curve (AUC) of the receiver operating characteristics (ROC) curve (Fielding and Bell, 1997; Phillips et al., 2006; Wiley et al., 2003). A variable's importance was measured as: 1) the permutation importance which compares the drop in training AUC if training and background data are permuted, and 2) jackknife of AUC for the test data when using only a single-variable model. Habitat suitability maps were predicted by applying Maxent models to all cells in the study region, using a logistic link function to yield habitat suitability index values (HSI) between zero and one (Phillips and Dudik, 2008).

We used comparable sets of variables for our overall, annual (5), and seasonal (4) models, with the exception that the burned – non-burned area variable was excluded from the seasonal model because our burned area data did not have sufficient temporal resolution.

## ***Results***

Distribution of Saiga antelope locations was uneven. Most of the locations were obtained from the northern part of the study area on the boundary between Stepnoi Nature Preserve and Chernye Zemli Nature Reserve. The terrain ruggedness index did not highlight areas of pronounced local changes in elevation. Overall, NW part of the study area had higher absolute heights than SE part. Most of the artificial water sources were located in the Nature Preserve with the exception of 5 of them, located to the north of the study area and in the unprotected island between Stepnoi and Chernye Zemli. Major part of the Nature Preserve was within 5000 meters from particular watering point. Distance to water sources in the Nature Reserve was usually higher than 5000 meters. In accordance with the allowed activities protection status, all farms

were also located along the eastern part of Stepnoi Nature Preserve and outside of it, but not in Chernye Zemli Nature Reserve. Most of the study area was burned at least once with the exception of the NE part where sheep numbers were still high and SE part where area burned only 1-2 times. Area that burned less was dominated by communities with higher share of shrubs and annuals. Frequently burned areas were located in the western part of the study area and corresponded with perennial and annual grasslands. Area under shrub-dominated communities in the Nature Reserve was negligible (Figure 29).

### **Overall variable importance**

Our overall habitat suitability model included ten predictors (cumulative burned area and burning frequency, distances to farms and water sources, vegetation in 1998 and 2007, mean and standard deviation NDVI and finally, absolute height and terrain ruggedness index). The overall model had a test AUC of 0.77 (range across the 5 replicate runs: 0.71–0.807) with a mean standard errors of 0.01. Habitat selection was mainly determined by five predictors: burning frequency, distance to farms, distance to water sources, standard deviation of NDVI, absolute height and total burned area, accounting for relative gain contributions of 36.6%, 24.4%, 18.6%, 5.9%, 3.5%, 3.4% respectively (combined, 92.4%) (Table 14).

### **Overall habitat distribution**

Saiga antelope habitat suitability was highest in the central part of the study area on the border between Stepnoi Nature Preserve and Chernye Zemli Nature Reserve. Saiga antelope habitat with an HSI value larger 0.5 covered about 20% of the study area (~454 km<sup>2</sup>), most of which occurred in one, contiguous patch. Habitat with an HSI value larger 0.6 covered only 7% of the study area (168 km<sup>2</sup>), and areas with an HSI value larger 0.7 were small, occupying about 2% of



the study area (Table 17). More suitable habitat ( $HSI > 0.6$ ) formed three separate patches; two of these patches were located in the northern part of the study area, where spring calving grounds are located, and the remaining one was located in to the south of this area.

### **Relationships between variables and habitat suitability in the overall model**

The overall model did not find substantial differences in habitat suitability between burned and unburned areas, and vegetation type in 1998 and 2007. For continuous variables, habitat suitability increased for areas that burned more often and areas with higher elevations. Saiga preferred also areas away from farms. Habitat suitability increased somewhat with increases in mean NDVI. Habitat suitability was negatively related to increasing distance to water sources. The standard deviation of NDVI, and terrain ruggedness had no apparent association with habitat suitability (Figure 34).

### **Annual and seasonal variability in variable importance**

As in the global model, the most important variables in the yearly and seasonal models for 2003-2007 were distance to water sources and burning frequency. These two variables combined contributed 60% to the overall model. The order of variable importance differed slightly among our two different metrics of variable importance (AUC on test data). Burning frequency was always first, elevation second, and distance to water sources and farms were the third and fourth most important variables (Table 16). Generally, the impact of vegetation-related variables, both NDVI-based and those derived from vegetation classifications, was lower than burning and distance variables. The binary burned / not-burned variable also had very high permutation importance in 2004.

## **Annual and seasonal variability in habitat distribution**

Spatially, we did not find substantial differences in the distribution of suitable habitat among years (Figure 31). Changes of more than 20% of HSI in a single year compared to HSI for all years were rare and located primarily in Chernye Zemli Nature Reserve (Figure 32). The distribution of suitable habitat predicted by the global model was similar to the distribution predicted by the yearly models. The total area covered by suitable habitat ( $HSI > 0.5$ ) was also comparable among years, with the exception of 2007, when it decreased from the average 20% of the total area for all years to about 10% (Table 17, Figure 32).

There were, however, some seasonal differences in the distributions of habitat suitability. In winter, suitable habitat was more likely to be found in the NW-corner of the study area, whereas in spring and summer suitable habitat was concentrated on the calving grounds in the north. In fall, suitable habitat was more dispersed. Overall though, these seasonal changes were not marked, and a general NW-SE gradient of habitat suitability was retained throughout the year (Figure 33). Areas covered by habitat seasonally with HSI values larger than 0.5 and 0.6 were comparable to the annual distributions, though somewhat smaller. Areas with HSI larger than 0.5 were least extensive in spring (14% of the study area) and most extensive in fall (20% of the study area) (Table 17).

## ***Discussion***

Our study represents one of the first attempts to understand habitat selection of Saiga antelope in the last remaining European population in Kalmykia (Russia) at the scale of the protected range. This is a constrained population that no longer migrates long distances. Our results indicate a higher importance of human-related factors, such as distance to water sources and farms, than

natural ones, such as vegetation, in determining Saiga antelope distributions. Patterns of suitable habitat among years and seasons were rather consistent. Saiga antelopes aggregated in the vicinity of the calving grounds even outside of the calving season. Most probably, the higher presence of Saiga antelope in this specific location is explained by a combination of factors: presence of water sources, it is the traditional area of the rut, and of calving (Arylova 2008), the area was predominantly flat compared to other parts of the study area, systematic fires improve visibility and forage, and there is little disturbance from vehicles and livestock.

We found a surprisingly strong influence of burning frequency in explaining habitat selection by Saiga antelope. Frequently burned areas might be more beneficial for Saiga antelope from both a forage and disturbance standpoint. Firstly, the areas that burn more frequently might have a specific vegetation composition favored by Saiga antelope. Grasses resprout rapidly after fire, even if there no rain and ungulates are shown to react on it by higher densities after fire (Archibald and Bond 2004; Everson and Everson 1987). The effect of burning on diet quality can persist for several years and be different depending on the season (Hobbs and Spowart 1984). Secondly, burning is a good predictor of the openness of the territory - areas that burned more often are characterized by improved visibility and passability for Saiga antelope (A. Khludnev, personal communication). Presence of continuous unburned vegetation cover made it hard for Saiga antelope to gain speed, and introduced a high level of noise from wind moving through plants, decreasing their ability to see and hear predators and poachers. We caution however, that it is possible that there was a reverse cause-effect relationship in which there is more burning because of saiga presence (i.e. fires set by poachers to aid hunting), but there are as yet no data to confirm this and there are many fires in the areas where Saiga are not present as well.

The low importance of vegetation in characterizing Saiga habitat suitability was certainly surprising. Soviet era studies and current research indicates that diet preferences are strongly seasonal and Saiga selected vegetation types that had greener biomass (e.g., shrub-dominated communities in the fall and winter and grasslands during spring-summer period). Recent studies on Saiga antelope diet indicate an avoidance of *Stipa* sp. altogether (Larionov et al. 2008). This is of concern, because of the continuing transition of the region to a greater dominance of this vegetation community (Abaturov 2007; Dubinin et al. 2010a). Fine-scale studies of Saiga antelope habitat selection in Kazakhstan emphasize the importance of green biomass measured by NDVI, especially during migration and calving (Leimgruber et al. 2001; Mueller et al. 2008; Singh et al. 2010b). However, our results do not support these finding - according to our models, neither greenness as measured by NDVI nor vegetation types were particularly strong predictors of suitable habitat, compared to our other variables. The lack of importance of vegetation variables indicated that the selection for different vegetation communities might be happening at finer scales, which were not captured by either our occurrence data or environmental layers, especially after generalization to 250-m pixels.

Though our study area was dominated by protected areas, the effect of human presence was still apparent in our models via the decreasing habitat suitability closer to farms. While we do not suggest that farmers themselves poach, farms are connected to transportation infrastructure which may make it easier for poachers to enter areas where Saiga antelope are present. Unfortunately, the sampling design and the use of off-road motorcycles by poachers impede modeling poaching pressure more directly. Finally, the very strong effect of distance to water sources highlights the aridity of the climate in the study area and the dependence of Saiga antelope on available water during the summer and even in spring and fall. Currently the area of

Chernie Zemli Nature Reserve does not have any artificial water sources and that may make the Stepnoi Nature Preserve more attractive to Saiga antelope. Habitat suitability in Chernie Zemli Nature Reserve may be increased through creation of more artificial water sources.

The occurrence dataset that we used is the only existing consistent dataset for our study area. However, this dataset suffered from a number of problems that could have potentially affected the results of our analysis. Firstly, the sampling effort does not follow specific protocol and oversampling in certain areas is likely a problem. Secondly, the locations do not represent the exact locations of the animals. This is why we aggregated the occurrence data (and all our predictor variables) to 250-m grid cells. Nevertheless, our training and test data were not truly independent and the resulting AUC of the models might be inflated due to autocorrelation in the residuals (Veloz 2009).

The clear patterns that we observed allowed us to make some management recommendations despite the limitations of the source data. We suggest that the key management implications for Saiga antelope resulting from this study are: habitat will be most effectively improved by optimizing the protected areas network, implementation of targeted anti-poaching activities, and enhanced monitoring. Because optimization of the network through removal of active farms from the management area (*Zakaznik*) is not feasible, we recommended building more water wells in the Nature Reserve where there are no farms, and no water sources either. Expanding the protected areas further to the north and center of our study area would also be helpful to fully capture currently unprotected portions of suitable habitat. Anti-poaching activities should be better focused on the area that we identified as suitable habitat. And last but not least, better wolf and saiga monitoring programs need to be launched to obtain more solid data on population dynamics and predation pressure. Given the ecological importance of fire, we

do not advise fire suppression, but rather suggest use of prescribed burning to ensure that areas burn at least once every two years.

The dramatic decline of Saiga antelope since the 1980s was concomitant with strong environmental changes, and especially a rapid increase in the area burned each year, and changes in vegetation types. This led to the hypothesis that habitat changes were the primary reason for Saiga antelope population collapse, but our habitat selection results do not support this hypothesis. Currently, the avoidance of disturbed areas by Saiga antelope, represented by poaching and predation, appears to be more important for the population than the presence of good foraging grounds. One reason for the lack of a clear relationship between vegetation type and saiga habitat quality might be that overall low densities of Saiga places population numbers far below their carrying capacity, and forage quality may therefore not be a limiting factor. However, the ongoing change of the area to more grass-dominated communities with impaired palatability for Saiga antelope should be carefully monitored to ensure that saiga prosper here in the longer term.

## ***Acknowledgements***

We gratefully acknowledge support for this research by the NASA Land-Cover and Land-Use Change Program (NNG05GA74G), by a NASA Earth System Science Dissertation Fellowship for M. Dubinin (07-Earth07F-0053), by the U.S. Fish and Wildlife Service, by a grant from the National Geographic Society (8360-07) and by Fundamental Research Program “Russia’s Biological Resources: assessment and monitoring fundamentals”, (Project PII.10 by the Department of Biology Sciences, Russian Academy of Sciences). We are also grateful to the

staff of Stepnoi Nature Preserve and Chernye Zemli Nature Reserve for collecting occurrence data. We thank T. Kuemmerle for help with Maxent and A. Estes for editing.

## ***Bibliography***

- Abaturov, B.D. (2007). The population of saiga antelopes in Russia and the problems of its preservation. *Herald of the Russian Academy of Sciences*, 77, 462-469
- Archibald, S., & Bond, W.J. (2004). Grazer movements: spatial and temporal responses to burning in a tall-grass African savanna. *International Journal of Wildland Fire*, 13, 377-385
- Arylova, N.Y. (2008). Reproduction features of the saiga antelope (*Saiga tatarica tatarica* L.) in Republic of Kalmykia in the low numbers' conditions. *Povolzhskii Ekologicheskii Zhurnal*, 2, 136-141
- Bannikov, A.G. (1967). The biology of the Saiga. *Israel Program for Scientific Translations, Jerusalem*, 1-252
- Bekenov, A.B., Grachev, I.A., & Milner-Gulland, E.J. (1998). The ecology and management of the Saiga antelope in Kazakhstan. *Mammal Review*, 28, 1-52
- Berger, J., Young, J.K., & Berger, K.M. (2008). Protecting Migration Corridors: Challenges and Optimism for Mongolian Saiga. *PLoS Biology*, 6, 1365-1367
- Bhatnagar, Y.V., Ahmad, R., Kyarong, S.S., Ranjitsinh, M.K., Seth, C.M., Lone, I.A., Easa, P.S., Kaul, R., & Raghunath, R. (2009). Endangered markhor *Capra falconeri* in India: through war and insurgency. *Oryx*, 43, 407-411
- Bowman, D., Zhang, Y., Walsh, A., & Williams, R.J. (2003). Experimental comparison of four remote sensing techniques to map tropical savanna fire-scars using Landsat-TM imagery. *International Journal of Wildland Fire*, 12, 341-348
- Chan, S., Maksimuk, A., & Zhirnov, L. (1995). From steppe to store: the trade in saiga antelope horn. *Trafic International*, 46



- Dubinin, M., Lushchekina, A., & Radeloff, V.C. (2010a). Fire and past landuse-driven vegetation change in semi-desert environments of Southern Russia (in preparation)
- Dubinin, M., Lushchekina, A., & Radeloff, V.C. (2010b). Relationships between climate, livestock, and fire in the arid grasslands of southern Russia (in review). *Ecosystems*
- Dubinin, M., Potapov, P., Lushchekina, A., & Radeloff, V.C. (2010c). Reconstructing long time series of burned areas in arid grasslands of southern Russia by satellite remote sensing. *Remote Sensing of Environment*, 114, 1638-1648
- Dudley, J.P., Ginsberg, J.R., Plumptre, A.J., Hart, J.A., & Campos, L.C. (2002). Effects of war and civil strife on wildlife and wildlife habitats. *Conservation Biology*, 16, 319-329
- Ehrlich, P.R., & Ehrlich, A.H. (1981). *Extinction: The Causes and Consequences of the Disappearance of Species*. New York: Ballantine Books
- Elith, J., Graham, C.H., Anderson, R.P., Dudik, M., Ferrier, S., Guisan, A., Hijmans, R.J., Huettmann, F., Leathwick, J.R., Lehmann, A., Li, J., Lohmann, L.G., Loiselle, B.A., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J.M., Peterson, A.T., Phillips, S.J., Richardson, K., Scachetti-Pereira, R., Schapire, R.E., Soberon, J., Williams, S., Wisz, M.S., & Zimmermann, N.E. (2006). Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, 29, 129-151
- Everson, C.S., & Everson, T.M. (1987). Factors Affecting the Timing of Grassland Regrowth after Fire in the Montane Grasslands of Natal South Africa. *South African Forestry Journal*, 47-52
- Golub, V.B. (1994). The desert vegetation communities of the Lower Volga Valley. *Feddes Repertorium*, 105, 499-515

- Hobbs, N.T., & Spowart, R.A. (1984). Effects of Prescribed Fire on Nutrition of Mountain Sheep and Mule Deer During Winter and Spring. *Journal of Wildlife Management*, 48, 551-560
- Karimova, T. (2002). What is needed for saiga conservation (results of specialists survey)? *Zapovednoye Delo*, 10, 98-107 [In Russian]
- Kuhl, A., Balinova, N., Bykova, E., Arylov, Y.N., Esipov, A., Lushchekina, A.A., & Milner-Gulland, E.J. (2009a). The role of saiga poaching in rural communities: Linkages between attitudes, socio-economic circumstances and behaviour. *Biological Conservation*, 142, 1442-1449
- Kuhl, A., Mysterud, A., Grachev, I.A., Bekenov, A.B., Ubushaev, B.S., Lushchekina, A.A., & Milner-Gulland, E.J. (2009b). Monitoring population productivity in the saiga antelope. *Animal Conservation*, 12, 355-363
- Kurinova, Z.S., & Belousova, Z.N. (1989). *Effective use of forage resources*. Elista: Kalmyk book publishing
- Larionov, K.O., Dzhapova, R.R., Rozenfeld, S.L., & Abaturov, B.D. (2008). Feeding of Saiga (Saiga Tatarica) on Pastures of Kalmykian Chernye Zemli under Plant Successions and Steppification. *Zoologicheskyy Zhurnal*, 87, 1259-1269
- Leimgruber, P., McShea, W.J., Brookes, C.J., Bolor-Erdene, L., Wemmer, C., & Larson, C. (2001). Spatial patterns in relative primary productivity and gazelle migration in the Eastern Steppes of Mongolia. *Biological Conservation*, 102, 205-212
- Milner-Gulland, E.J., Bekenov, A.B., & Grachov, Y.A. (1995). The real threat to saiga antelopes. *Nature*, 377, 488-489
- Milner-Gulland, E.J., & Bennett, E.L. (2003). Wild meat: the bigger picture. *Trends in Ecology & Evolution*, 18, 351-357

- Milner-Gulland, E.J., Bukreeva, O.M., Coulson, T., Lushchekina, A.A., Kholodova, M.V., Bekenov, A.B., & Grachev, I.A. (2003). Reproductive collapse in saiga antelope harems. *Nature*, 422, 135
- Milner-Gulland, E.J., Kholodova, M.V., B., B.A., Bukreeva, O.M., Grachev, I.A., Amgalan, L., & Lushchekina, A.A. (2001). Dramatic declines in saiga antelope populations. *Oryx*, 35, 340-345
- Morrison, J.C., Sechrest, W., Dinerstein, E., Wilcove, D.S., & Lamoreux, J.F. (2007). Persistence of large mammal faunas as indicators of global human impacts. *Journal of Mammalogy*, 88, 1363-1380
- Mueller, T., Olson, K.A., Fuller, T.K., Schaller, G.B., Murray, M.G., & Leimgruber, P. (2008). In search of forage: predicting dynamic habitats of Mongolian gazelles using satellite-based estimates of vegetation productivity. *Journal of Applied Ecology*, 45, 649-658
- Phillips, S.J., Anderson, R.P., & Schapire, R.E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190, 231-259
- Phillips, S.J., Dudik, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J., & Ferrier, S. (2009). Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecological Applications*, 19, 181-197
- Riley, S.J., DeGloria, S.D., & Elliot, R. (1999). A terrain ruggedness index that quantifies topographic heterogeneity. *Intermountain Journal of Sciences*, 5
- Robinson, S. (2000). Pastoralism and land degradation in Kazakhstan. In, *PhD thesis*: University of Warwick, UK
- Simberloff, D. (1984). Mass Extinction and the Destruction of Moist Tropical Forests. *Zhurnal Obshchei Biologii*, 45, 767-778

- Singh, N.J., Grachev, I.A., Bekenov, A.B., & Milner-Gulland, E.J. (2010a). Saiga antelope calving site selection is increasingly driven by human disturbance. *Biological Conservation*, *143*, 1770-1779
- Singh, N.J., Grachev, I.A., Bekenov, A.B., & Milner-Gulland, E.J. (2010b). Tracking greenery across a latitudinal gradient in central Asia – the migration of the saiga antelope. *Diversity and Distributions*, *16*, 663-675
- Sodhi, N.S., Koh, L.P., Brook, B.W., & Ng, P.K.L. (2004). Southeast Asian biodiversity: an impending disaster. *Trends in Ecology & Evolution*, *19*, 654-660
- Sokolov, V.E., & Zhirnov, L.V. (Eds.) (1998). *Saigak: filogeniya, sistematika, ekologiya, okhrana i ispol'zovanie [The saiga antelope: phylogeny, systematics, ecology, conservation, and use]*. Moscow: Rosselkhozacademiya
- Veloz, S.D. (2009). Spatially autocorrelated sampling falsely inflates measures of accuracy for presence-only niche models. *Journal of Biogeography*, *36*, 2290-2299
- Walter, H., & Box, E. (1983). Overview of Eurasian continental deserts and semideserts. *Ecosystems of the World* (pp. 3-269). Amsterdam Elsevier
- Wilkie, D., Shaw, E., Rotberg, F., Morelli, G., & Auzel, P. (2000). Roads, development, and conservation in the Congo basin. *Conservation Biology*, *14*, 1614-1622

## Tables

Table 13. Predictors used to model suitable habitat and determine its drivers and variability in Southern Russia.

Variable	Type	Range	Source	Models
Burned area	Categorical	0-1	MODIS	annual, overall
Burned area	Continuous	0-6	MODIS	overall, annual, seasonal
Vegetation in 2007	Categorical	1-9 (see text for classes description)	Landsat	overall, annual, seasonal
Vegetation in 1998	Categorical	1-9 (see text for classes description)	Landsat	overall, annual, seasonal
Elevation	Continuous	-37 – -9 (m)	SRTM	overall, annual, seasonal
Terrain ruggedness	Continuous	0 – 20	SRTM	overall, annual, seasonal
Distance to farms	Continuous	0 – 25,500 (m)	Topographic maps	overall, annual, seasonal
Mean NDVI	Continuous	0 – 10,000	MOD13A1	overall, annual, seasonal
Standard deviation NDVI	Continuous	0 – 10,000	MOD13A1	overall, annual, seasonal
Distance to water sources	Continuous	0 – 27,000 (m)	Topographic maps	overall, annual, seasonal

Table 14. Amount of data used for model construction and resulting performance of the models.

SD AUC represents average standard deviation across 5 replications, while each SD AUC is calculated by folding and using remaining folds for cross-validation, repeating for every replication.

	250-m				
	grid		Test		
	Occurrence	occurrence	Training	data	SD
	points	points	data AUC	AUC	AUC
2003	120	104	0.84	0.81	0.04
2004	260	133	0.81	0.86	0.03
2005	573	196	0.78	0.77	0.03
2006	428	251	0.77	0.71	0.03
2007	482	277	0.86	0.85	0.02
Winter	614	313	0.81	0.77	0.02
Spring	431	225	0.83	0.79	0.03
Summer	411	213	0.81	0.75	0.03
Fall	523	317	0.81	0.77	0.02
Overall	1979	851	0.77	0.77	0.01

Table 15. Variable contribution measured as permutation importance, percentages, averaged over 5 runs: a) overall and annual models, b) seasonal models

a)

	Overall	2003	2004	2005	2006	2007	sum	mean
burning frequency	27.8	35	14	35	36	24	143	29
burned-non burned	0	0	25	0	5	1	32	6
distance to water	35	21	24	13	15	51	123	25
distance to farm	12	6	12	6	10	8	42	8
elevation	7	8	8	27	5	6	53	11
terrain ruggedness	1	1	4	6	3	2	15	3
vegetation type in 1998	1	1	4	2	3	0	11	2
vegetation type in 2007	0	5	5	8	17	5	40	8
mean NDVI	4	13	1	1	6	0	21	4
standard deviation								
NDVI	10	9	3	2	2	3	19	4

b)

	Winter	Spring	Summer	Fall	sum	mean
burning frequency	44	29	20	24	116	29
distance to water	11	38	34	38	121	30
distance to farm	8	10	13	11	42	10
elevation	16	7	11	5	38	10
terrain ruggedness	4	2	5	1	13	3

vegetation type in 1998	1	2	2	2	7	2
vegetation type in 2007	2	6	13	3	24	6
burned-non burned	7	3	1	6	17	4
mean NDVI	8	3	2	10	23	6
standard deviation						
NDVI	44	29	20	24	116	29

---



Table 16. Results of jackknife test for variable importance as AUC on test data sorted by sum, as percentages, averaged over 5 runs: a) overall and annual models, b) seasonal models

a)

	Overall	2003	2004	2005	2006	2007	sum	mean
burning frequency	70	69	66	74	69	71	348	70
burned-non burned	63	50	70	48	66	53	288	58
distance to water	62	66	59	54	58	74	310	62
distance to farms	62	60	55	57	59	66	297	59
elevation	63	63	65	70	63	65	326	65
terrain ruggedness	53	51	49	52	50	52	254	51
vegetation type in 2007	62	50	54	53	55	58	270	54
vegetation type in 1998	54	58	65	59	64	63	310	62
mean NDVI	53	58	62	56	50	55	282	56
standard deviation NDVI	58	66	66	55	56	57	300	60

b)

	Winter	Spring	Summer	Fall	sum	mean
burning frequency	72	69	68	71	280	70
distance to water	54	56	57	55	222	56
distance to farms	64	64	60	62	251	63
elevation	58	64	60	60	242	61
terrain ruggedness	57	67	66	65	256	64
burned-non burned	71	64	62	64	260	65

vegetation type in 1998	52	51	48	51	201	50
vegetation type in 2007	58	47	51	57	213	53
mean NDVI	63	61	65	59	248	62
standard deviation NDVI	72	69	68	71	280	70

---

Table 17. Amount of suitable and highly suitable habitat and its change across years and seasons in percents of total study area.

	>50%	>60%	>70%
2003	18%	8%	2%
2004	16%	10%	3%
2005	22%	9%	2%
2006	23%	8%	2%
2007	10%	5%	2%
Winter	17%	7%	2%
Spring	14%	6%	2%
Summer	16%	6%	2%
Fall	20%	7%	2%
Overall	22%	8%	2%

## Figures

Figure 27. Dynamics of livestock and Saiga antelope populations from 1952 to 2008 in Kalmykia.




Figure 28. Study area location in southern Russia. Dashed lines in the study area represent protected areas (left - Stepnoi Nature Preserve and right - Chernye Zemli Nature Reserve), hatched area with black outline – study area, thick grey line – administrative boundary of Republic of Kalmykia.



Figure 29. Distribution of selected variables. Clockwise from the top left: elevation (meters), burning frequency (number of times a location burned), distance to water sources (thousands of meters), distance to farms (thousands of meters), vegetation communities in 2007 (for legend see Methods section), mean total NDVI (index value). Dashed lines represent boundaries of protected areas.




Figure 30. Habitat suitability map using overall dataset showing the point-wise mean. Blue represents preferred habitat, brown non-preferred, white missing data. Violet dots represent Saiga antelope locations used as training points, green - locations used for validation. Dots represent one of the replications and provided as an example. Dashed lines represent boundaries of protected areas.




Figure 31. Annual distribution of suitable habitat, averages over 5 replications are shown.






Figure 32. Difference of the habitat suitability index for a given year minus the habitat suitability index for all years 2003 – 2007).




Figure 33. Seasonal distribution of suitable habitat, averages over 5 replications are shown.




Figure 34. Response curves showing relationship between the particular variable and logistic output (probability of presence) according to the global model. Gray lines indicate minimum and maximum from 5 replications and thick black line represents a mean. See Table 1 for explanations of units of x-axis.

